Thioautotrophic Bacterial Endosymbionts Are Degraded by Enzymatic Digestion During Starvation: Case Study of Two Lucinids *Codakia orbicularis* and *C. orbiculata*.

Sten König, Hervé Le Guyader, Olivier Gros

► To cite this version:

Sten König, Hervé Le Guyader, Olivier Gros. Thioautotrophic Bacterial Endosymbionts Are Degraded by Enzymatic Digestion During Starvation: Case Study of Two Lucinids *Codakia orbicularis* and *C. orbiculata*.. Microscopy Research and Technique, 2015, 78 (2), pp.173-179 10.1002/jemt.22458 . hal-01206367

HAL Id: hal-01206367

<https://hal.univ-antilles.fr/hal-01206367>

Submitted on 8 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Thioautotrophic bacterial endosymbionts are degraded by lysosomal host digestion during starvation: case study of two lucinids -*Codakia orbicularis* and *Codakia orbiculata*.

Sten König¹, Hervé Le Guyader² and Olivier Gros^{1,3}

¹ *Institut de Biologie Paris-Seine, UMR 7138 - Evolution Paris-Seine, Equipe Biologie de la Mangrove. Université des Antilles et de la Guyane, UFR des Sciences Exactes et Naturelles, Département de Biologie, BP 592. 97159 Pointe-à-Pitre cedex, Guadeloupe, France*

² *Sorbonne Universités Paris VI, Institut de Biologie Paris-Seine, UMR 7138 - Evolution Paris-Seine, Equipe Phylogénie, Anatomie, Evolution. C.N.R.S, Institut de Biologie Paris-Seine, UMR 7138 - Evolution Paris-Seine, Equipe Biologie de la Mangrove.*

³ *C₃MAG, UFR des Sciences Exactes et Naturelles, Université des Antilles et de la Guyane, BP 592 - 97159 Pointe-à-Pitre, Guadeloupe, French West Indies.*

Sten König: *sten.koenig@gmail.com*

Olivier Gros: *olivier.gros@univ-ag.fr*

Hervé Le Guyader: *herve.Le_Guyader@upmc.fr*

Key words: chemoautotrophicsymbiosis, cytochemistry, EDXS analysis, TEM, bacterial degradation

running title: Endosymbiont digestion in the bivalves *Codakia*.

contact: Sten König
sten.koenig@gmail.com
0049-(0)3834-864915
University Greifswald
Institut of Pharmacy
Friedrich-Ludwig-Jahn Straße 17
17489 Greifswald, Germany

Abstract

The Caribbean bivalves *Codakia orbicularis* (Linné, 1758) and *C. orbiculata* (Montagu, 1808) live in seagrass beds of *Thalassia testudinum* and harbor intracellular sulfur-oxidizing Gamma-proteobacteria. These bacterial symbionts fix CO₂ via the Calvin Benson Cycle and provide organic compounds to the bivalve. In case of starvation, no reduced sulfur compounds and no organic particle food are available; the symbionts could be considered as the sole nutrient source by the host. In previous studies it was shown that the intracellular bacterial population decreased considerably during starvation and that the bacteria were not released by the bivalves (Brissac et al., 2009). In the present study, the activity of two lysosomal marker enzymes (acid phosphatase and arylsulfatase) was detected using cytochemical experiments coupled with EDX-TEM (Energy Dispersive X-ray-Transmission Electron Microscopy) during sulfide and organic particle starvation. The degradation of bacterial endosymbionts began after two weeks of starvation in *C. orbiculata* and after three weeks in *C. orbicularis*. Such degradation process seems continuous through months and could be responsible for the disappearance of the bacterial endosymbionts within the gills during starvation. These data suggest that the host uses symbionts as a nutrient source to survive a hunger crisis. The carbon transfer from the symbionts to the host could be flexible and could consist in transfer of organic matter, "milking", under normal feeding conditions and digestion of the symbionts, "farming", under starved conditions.

Introduction

The association between chemoautotrophic bacterial symbionts and marine invertebrates was first discovered in the deep sea at hydrothermal vents (Felbeck, 1981). In shallow-water environments, similar symbiotic relationships were described involving bivalves (Felbeck, 1983; Reid, 1990), nematodes (Ott et al., 1982) and annelids (Cavanaugh et al., 1981; Felbeck, 1981). To date it is known that the entire bivalve family Lucinidae harbors chemoautotrophic Gamma-proteobacteria in specialized gill cells called bacteriocytes (Taylor et al., 1997). Many lucinid bivalves inhabit seagrass beds, where they dominate the sedimentary infauna (Reynolds et al., 2007). The seagrass ecosystem is a very productive system and is enriched in organic matter (Duarte et al., 1999). The anaerobic degradation of organic matter in seagrass bed sediments by sulfate reducers produces reduced sulfur compounds such as hydrogen sulfide (Jørgensen, 1982). Reduced sulfur compounds (herein called sulfides) are toxic to most plants and animals (Dorman et al., 2002). Lucinid bivalves live at the interface of oxic and anoxic sediment zones. They build burrows in sulfide-rich anaerobic sediments and are connected to the surface water, which is saturated with oxygen. The bacterial endosymbionts oxidize sulfides, produce organic matter through the Calvin Benson Cycle and supply these organic compounds to the host (Van Dover, 2000). In a direct carbon transfer called "milking", the symbiont permanently translocate organic compounds to the host soon after carbon fixation (Schweimanns and Felbeck, 1985; Streams et al., 1997; Van Dover, 2000). Therefore, the digestion system and filtering capacity of lucinids are reduced (Reid, 1990) compared to other bivalves without endosymbionts. Lucinids provide their endosymbionts with sulfide and oxygen through permanently water pumping, oxygen acts as terminal electron acceptor.

Codakia orbicularis (Linné, 1758) and *C. orbiculata* (Montagu, 1802) inhabit the shallow water seagrass bed of *Thalassium testudinum* in the Caribbean Sea. Both lucinids have been studied with regard to the interaction between host and symbionts in the last years, these systems represent an attractive model to investigate symbiosis. According to 16S rRNA sequence analysis, both *Codakia* species harbor the same and only one bacterial symbiont phylotype inside the gills (Durand et al., 1996). Cultivation of aposymbiotic juveniles of *C. orbicularis* in controlled conditions and in natural seagrass bed sediment demonstrate that the transmission of the gill-endosymbionts is environmental (Gros et al., 1996). Free living symbionts were detected in seagrass bed sediments (Gros et al., 1996; 2003). No gill-endosymbionts were detectable in either larvae or juveniles of *C. orbicularis*, suggesting that

the bivalve acquires symbiosis-competent bacteria once their gill tissue is sufficiently developed (Gros et al., 1998). *C. orbiculata* can acquire bacteria during their whole life by phagocytosis at the apical pole of the bacteriocytes (Elisabeth et al., 2012). The symbionts of both lucinids are not released by the host, so the intracellular life of the symbionts seems to be a dead end for the bacteria (Brissac et al., 2009). Caro et al. (2007) suggested that *C. orbicularis* could control bacterial growth to harvest bacterial biomass. Interestingly, it was shown that there are up to seven sub-populations of this symbiont phylotype within a single bivalve. These bacterial sub-populations differ in size, number of chromosome copies and content of sulfur granules (Caro et al., 2007, 2009).

The gills of lucinids are divided in three zones i) the ciliated zone ii) the intermediary zone, and iii) the lateral zone, which takes up most of the volume of the gills (Frenkiel et al., 1995). The ciliated and intermediary zones are free of symbiotic bacteria. The upper two thirds of the lateral zone are occupied by bacteriocytes (which harbor bacterial symbionts) and by intercalary cells. The lower third is mainly occupied by granule cells, considered to be important for storage and metabolic conversion of sulfur compounds (Frenkiel and Mouëza, 1995). In case of starvation the organization of the gills in *C. orbicularis* changes and the bacterial population decreases by about one third per month (Caro et al., 2009). After six months of starvation, no symbiont is detectable in the gills, the main part of which is then occupied by granule cells. The reduction of bacteriocyte's volume occurs parallel to the reduction of the number of bacterial symbionts (Caro et al., 2009). Due to the fact that no bacteria were detectable in these sea-water tanks used during lucinid starvation experiments, Brissac et al. (2009) concluded that there was no release of the bacterial symbionts. The decrease in size of the symbiont population could be either due to bacterial autolysis or to host-driven lysis, as a mean of obtaining nutrients from these bacteria. Indeed, bivalves with chemoautotrophic symbionts are known to digest and use symbionts as a nutrient source (Herry et al., 1989; Boetius and Felbeck, 1995; Fiala et al., 1994; Liberge et al., 2001). The direct digestion of the symbiont and the subsequent nutrient transfer to the host is called "farming" (Streams et al., 1997). Symbiont autolysis has been described in *Bathymodiolus azoricus* (Kádár et al., 2007). In this case the author hypothesized that the autolysis of the bacteria is part of the cell cycle and that a bacterial acid phosphatase is involved in the process of the autolysis. Acid phosphatase and arylsulfatase are common enzymes in gram-negative bacteria (Rogers, 1960; Chandramohan et al., 1974). In eukaryotes, acid phosphatase and arylsulfatase are lysosomal markers that have been detected in the Mollusca (Patel and Patel, 1985; Patel and Patel, 1985; Pipe and Moore, 1985; Liberge et al., 2001). Lysosomes are eukaryotic

membrane bound cell organelles with hydrolytic enzymes to chemically deteriorate polymers and bacteria. Acid phosphatase and arylsulfatase are two marker enzymes for lysosomal bodies (Löffler et al., 1984). Here, we document changes in acid phosphatase and arylsulfatase activity in the gills of *C. orbicularis* and *C. orbiculata* during host starvation. We suggest that the host can use this large amount of bacteria as nutrient source through lysosomal degradation.

Material and Methods

Sampling of bivalves

Adult individuals of *Codakia orbicularis* and *C. orbiculata*, were manually collected in *Thalassium testudinum* seagrass beds on "Îlet Cochon" (16°12'53"N; 61°32'05"W) in Guadeloupe (French West-Indies, Caribbean).

Laboratory maintenance of the bivalves

Lucinids (N=80 *C. orbiculata* and N=40 *C. orbicularis*) were kept in separate 50L plastic tanks with 0.22 µm filtered seawater at 26°C up to six months. The water was oxygenated with an aquarium air pump. No food (organic particles) or reduced sulfur compounds were added to the water throughout the starvation experiment. The water in the tanks was changed every seven days. At specific times, three bivalves were randomly sampled from the tanks for analysis; for *C. orbiculata*, sampling took place upon collection and then every two weeks through the experiment (i.e 5 months) while for *C. orbicularis*, sampling took place three weeks after initial field collection then every two weeks.

Cytochemistry

The gills of the bivalves were dissected and fixed (2% glutaraldehyde, 1% paraformaldehyde in 0.1 M cacodylate adjusted to 900 mOsm,) for 2h at 4°C. The fixed gills were washed overnight at 4°C in 0.2M sodium cacodylate buffer with an adjusted osmolarity at 1800 mOsm with NaCl and Ca₂Cl. The gills were sliced in sodium cacodylate buffer into 100 µm sections using a Vibratome OCT Slicer at 4°C. Only one enzyme activity can be detected per gill section, therefore for each enzyme reaction three replicate sections were prepared. Arylsulfatase activity was detected according to a method from Hopsu-Havu et al. (1967 cited in, Narbaitz, 1995). Briefly, the 100µm sections were transferred directly to the enzyme reaction solution (60 mM sodium acetate, 15 mM acetic acid, 8 mg/ml sulfate p-nitrocatechol, 60 mM barium chloride) and incubated for 45 min at 37°C. After the incubation, the samples were washed in cacodylate buffer in several changes of sodium cacodylate buffer overnight.

Acid phosphatase was detected by a method from Barka and Anderson (Barka et al., 1962). The sections were pre-incubated for 10 to 20 min at 37°C in 50 mM acetate buffer with 15 mM acetic acid. The enzymatic reaction took place in the pre-incubation buffer containing an additional 0.1 M sodium β -glycerophosphate substrate, saturated with lead-(II)-nitrate. The sections were incubated for 45 min at 37°C. After incubation the sections were washed in acetate buffer over a period of 60 min with several changes of the washing buffer. For both acid phosphatase and arylsulfatase assays, a precipitate containing the heavy metal was formed as a result of enzyme activity during the incubation stage. For both enzymes a negative control was prepared therefore the 100 μ m gill sections were incubated in the particular enzyme reaction solution without the enzyme substrate but with the heavy metal. Following incubations, 100 μ m sections were processed for embedding. Samples were dehydrated through a graded acetone series and embedded in Epon-araldite resin according to Glauert (1991). Ultra-thin (100 nm) sections were made using an ultracut E microtome and two grids containing 4-5 sections were observed with a Tecnai G2 TEM at 200 kv. The barium or lead precipitates were detected using a 30mm² EDX detector. The EDXS analysis corresponds to an analytical method used for elemental characterization. Micrographs presented in this study are representative of the three replicate bivalves examined at each sampling time during starvation.

Results

During starvation, the gills of the bivalves changed visibly from thick and beige to thin and dark brown. The mortality of the bivalves in the sterile sea water tanks was low: around one to five percent per week. Although the lethality was higher during the first three weeks compared to the later five months of starvation. On transmission electron micrographs, positive enzymatic reactions appeared as black areas, resulting from the reaction of acid phosphatase and arylsulfatase with lead and barium, respectively.

In freshly collected *C. orbiculata*, gill filaments showed weak positive reaction: only few black areas were visible in a couple of bacteriocytes (fig 1 A/B) while most of the bacteriocytes throughout the gill filaments are free of lysosomal enzyme activity (data not shown). This data suggests that adult individuals in their natural habitat can digest symbionts, probably in the frame of resorption of dead bacteria. After two weeks of starvation, the acid phosphatase activity was detected close to symbionts, while other gill cells were free of enzyme activity (fig

1 C/D). Likewise, the activity of the arylsulfatase increased during starvation time, after two months most of the symbionts inside the bacteriocytes are affected by an arylsulfatase activity (fig 1 E). After the beginning of bacterial degradation in the gills, the lysosomal activity was observed in all bacteriocytes of the lateral zone. It seems that it is a global process in the bacteriocytes of the gills in case of starvation. The further the course of starvation in both bivalves went, the amount of black areas increased until most of the bacteria were degraded. Nevertheless, for long period of starvation (5 months and more), the lysosomal activity globally decreases as less bacteriocytes are present through the lateral zone which now contains mostly granule cells. However, in the few remaining bacteriocytes which contain only few bacterial symbionts, there is still a lysosomal digestion occurring. No black areas were observed in granule cells and intercalary cells.

For *C. orbicularis* the story looked similar. After three weeks of starvation, no positive signal was detected for arylsulfatase and acid phosphatase in the gills of *C. orbicularis*, indicating very weak to no lysosomal activity (fig 2 A/B). In *C. orbicularis*, both enzyme activities were detected after six weeks of starvation (fig 2 C/D/E). The longer the starvation period, the more extensive was the area corresponding to the enzyme activity, until most symbionts seemed affected (data not shown). Similarly to *C. orbiculata*, the enzymatic activity in *C. orbicularis* increased during the first months of starvation, then progressively decreased after 4-5 months of starvation when the number of bacteriocytes and bacterial gill-endosymbionts have been strongly reduced through the lateral zone. The EDX analysis confirmed that the black areas contained barium and lead demonstrating that arylsulfatase and acid phosphatase enzymes were efficient (fig. 1F/2F). In the negative controls, no barium or lead was detectable (not shown).

Discussion and Conclusion

The present study uses cytochemical methods to explain how lucinid bivalves survive when no organic particulate food or sulfides (needed for symbiont thiotrophic metabolism) are available. The symbionts occupy around 34% of the gills of freshly collected *C. orbicularis* (Caro et al., 2009). During starvation, the bacterial population decreases by one third per month (Caro et al., 2009) while no released bacteria are detectable using CARD-FISH (Brissac et al., 2009). Lysosomes were rarely observed in freshly collected gills of *C. orbicularis* (Frenkiel et al., 1995). In our study in freshly collected *Codakia* bivalves, symbiont digestion was rarely observed. These data suggest that adult individuals in their natural habitat can digest symbionts, probably in the frame of resorption of dead bacteria and symbiont population controlling. In deep sea bivalves it was suggested that lysosomal

symbiont digestion is a way of controlling symbiont population (Fiala-Medioni et al., 1990). During starvation, lysosomes in the gills in both bivalves become larger and more abundant (Caro et al., 2009). Moreover, Elisabeth et al. (2012) described a massive decrease of the area occupied by bacteriocytes in *C. orbiculata* during starvation. *C. orbiculata* can acquire its symbionts in all life stages, from a free living stock of symbionts in the seagrass sediment (Gros et al., 2003) probably through phagocytosis at the apical pole of bacteriocytes (Gros et al., 2012). Inside the gills, the proliferation of the bacteria is inhibited by an unknown mechanism (Caro et al., 2007). The molecular mechanisms that regulate symbiont proliferation likely involve complex symbiont-host signalling processes (Stewart et al., 2006). In the laboratory, it is possible to maintain adult bivalves under starvation conditions for up to six months, and minimize the number of symbionts inside the gills. During our experiments, the appearance of the lucinid gills changed from thick and lightly beige in freshly collected bivalves (containing large symbiont population with a big amount of elemental sulfur) to thin and dark brown in starved bivalve individuals (containing less symbionts without elemental sulfur) (personal observations) (Johnson et al., 2001; Lechaire et al., 2008; Caro et al., 2009). We suggest that the host can use this large amount of bacteria as nutrient source through lysosomal degradation. Although their digestive tract is reduced (Reid, 1990) particle feeding is still part of the nutrient strategy of symbionts-bearing lucinids (Duplessis et al., 2004). During starvation, lysosomal enzyme activities were observed inside bacteriocytes close to, and within, the bacteria. We suggest that the host digests its bacterial endosymbionts during starvation to use the bacteria themselves as nutrients (i.e. farming). The trophic pathway has been described in other marine chemotrophic symbioses (Herry et al., 1989; Fiala et al., 1994; Boetius and Felbeck, 1995; Kádár et al., 2007). For example, the lucinid *Lucina pectinata*, digests its symbionts in large secondary lysosomes (Liberge et al., 2001). The enzymatic degradation of the symbionts in *Codakia* appeared different from that observed in *L. pectinata* as no large secondary lysosomes with a positive activity of the two lysosomal enzymes tested were observed during this study in *C. orbicularis* bacteriocytes. In *Codakia* individuals, the gill-endosymbionts seem to be individually digested compared to multiple bacteria digestion with large lysosomes in *L. pectinata*. It seems that *Codakia* is less adapted to symbiont digestion than *L. pectinata*.

Most of the enzyme activity was observed inside the symbionts indicating that the detected enzymes could be produced by symbionts themselves and be active even after their death. Symbionts could not find sulfides and/or enough nutrients inside the bacteriocytes due to the fact that stressed hosts preferred to sustain their own metabolism at the expense of its symbiotic population. So, once died and autolysed, the degraded products from the

symbionts could possibly become incorporated by host cells helping the hosts to survive starved conditions. Then, in better conditions, hosts will acquire new symbiosis-competent bacteria from its environment restoring a “normal” metabolic pathway. It remains uncertain whether lucinids actively lyse bacteria to obtain nutrients during periods of host starvation, or whether the symbionts, starved of their energy source, undergo autolysis (with possible nutritional benefits to the host). Further studies are necessary to investigate how bacterial populations are controlled by lucinid hosts under natural conditions using metabolomics or stable isotope analyses.

Acknowledgments

StenKönig was funded through a Ph.D. grant from the EU Marie Curie Initial Training Network (ITN) "Symbiomics". Finally, we would like to thank the three anonymous reviewers whose suggestions helped us to improve our manuscript.

References

- Barka, T., and Anderson, P.J. (1962). Histochemical methods for acid phosphatase using hexazonium pararosanilin as coupler. *J. Histochem. Cytochem.* 10, 741–753.
- Boetius, A., and Felbeck, H. (1995). Digestive enzymes in marine invertebrates from hydrothermal vents and other reducing environments. *Mar. Biol.* 122, 105–113.
- Brissac, T., Gros, O., and Mercot, H. (2009). Lack of endosymbiont release by two Lucinidae (Bivalvia) of the genus *Codakia*: consequences for symbiotic relationships. *FEMS Microbiol. Ecol.* 67, 261–267.
- Caro, A., Gros, O., Got, P., Wit, R.D., and Troussellier, M. (2007). Characterization of the population of the sulfur-oxidizing symbiont of *Codakia orbicularis* (Bivalvia, Lucinidae) by single-cell analyses. *Appl. Environ. Microbiol.* 73, 2101–2109.
- Caro, A., Got, P., Bouvy, M., Troussellier, M., and Gros, O. (2009). Effects of long-term starvation on a host bivalve *Codakia orbicularis* (Lucinidae) and its symbiont population. *Appl. Environ. Microbiol.* 75, 3304–3313.
- Cavanaugh, C.M., Gardiner, S.L., Jones, M.L., Jannasch, H.W., and Waterbury, J.B. (1981). Prokaryotic cells in the hydrothermal vent tube worm *Riftia pachyptila* Jones: possible chemoautotrophic symbionts. *Science* 213, 340–342.
- Chandramohan, D., Devendran, K., and Natarajan, R. (1974). Arylsulfatase activity in marine sediments. *Mar. Biol.* 27, 89–92.
- Dorman, D.C., Moulin, F.J.-M., McManus, B.E., Mahle, K.C., James, R.A., and Struve, M.F. (2002). Cytochrome oxidase inhibition induced by acute hydrogen sulfide inhalation: correlation with tissue sulfide concentrations in the rat brain, liver, lung, and nasal epithelium. *Toxicol. Sci.* 65, 18–25.

Duarte, C.M., and Chiscano, C.L. (1999). Seagrass biomass and production: a reassessment. *Aquat. Bot.* 65, 159–174.

Duplessis, M.R., Dufour, S.C., Blankenship, L.E., Felbeck, H., and Yayanos, A.A. (2004). Anatomical and experimental evidence for particulate feeding in *Lucinoma aequizonata* and *Parvilucina tenuisculpta* (Bivalvia: Lucinidae) from the Santa Barbara Basin. *Mar. Biol.* 145, pages number??.

Durand P., P., Gros, O., Frenkiel, L., and Prieur, D. (1996). Phylogenetic characterization of sulfur-oxidizing bacterial endosymbionts in three tropical Lucinidae by 16S rDNA sequence analysis. *Mol Mar Biol Biotechnol* 5, 37–42.

Elisabeth, N.H., Gustave, S.D.D., and Gros, O. (2012). Cell proliferation and apoptosis in gill filaments of the lucinid *Codakia orbiculata* (Montagu, 1808) (Mollusca: Bivalvia) during bacterial decolonization and recolonization. *Microsc. Res. Tech.* 75, 1136–1146.

Felbeck, H. (1981). Chemoautotrophic potential of the hydrothermal vent tube worm, *Riftia pachyptila* Jones (Vestimentifera). *Science* 213, 336–338.

Felbeck, H. (1983). Sulfide oxidation and carbon fixation by the gutless clam *Solemya reidi*: an animal-bacteria symbiosis. *J. Comp. Physiol.* 152, 3–11.

Fiala, M.A., Michalski, J.C., Jolles, J., Alonso, C., and Montreuil, J. (1994). Lysosomal and lysozyme activities in the gill of bivalves from deep hydrothermal vents. *C. R. Acad. Sci. III* 317, 239–244.

Fiala-Medioni, A., Felbeck, H., Childress, J.J., Fisher, C.R., and Vetter, R.D. (1990). Lysosomal resorption of bacterial symbionts in deep-sea bivalves. In *Endocytobiology IV: 4th International Colloquium on Endocytobiology and Symbiosis*, INSA-Villeurbanne (France), July 4-8, 1989, P. Nardon, V. Glaninazzi-Pearson, A.M. Grenier, L. Margulis, and D.C. Smith, eds. (Paris: Institut national de la recherche agronomique), pp. 335–338.

Frenkiel, L., and Mouëza, M. (1995). Gill ultrastructure and symbiotic bacteria in *Codakia orbicularis* (Bivalvia, Lucinidae). *Zoomorphology* 115, 51–61.

Glauert, A.M. (1991). Epoxy resins: an update on their selection and use. *Eur. Microsc. Anal.* 5, 15–20.

Gros, O., Darrasse, A., Durand, P., Frenkiel, L., and Mouëza, M. (1996). Environmental transmission of a sulfur-oxidizing bacterial gill endosymbiont in the tropical lucinid bivalve *Codakia orbicularis*. *Appl. Environ. Microbiol.* 62, 2324–2330.

Gros, O., Frenkiel, L., and Mouëza, M. (1998). Gill filament differentiation and experimental colonization by symbiotic bacteria in aposymbiotic juveniles of *Codakia orbicularis* (Bivalvia: Lucinidae). *Invertebr. Reprod. Amp Dev.* 34, 219–231.

Gros, O., Liberge, M., Heddi, A., Khatchadourian, C., and Felbeck, H. (2003). Detection of the free-living forms of sulfide-oxidizing gill endosymbionts in the lucinid habitat (*Thalassia testudinum* environment). *Appl. Environ. Microbiol.* 69, 6264–6267.

Gros, O., Elisabeth, N.H., Gustave, S.D.D., Caro, A., and Dubilier, N. (2012). Plasticity of symbiont acquisition throughout the life cycle of the shallow-water tropical lucinid *Codakia orbiculata* (Mollusca: Bivalvia): Symbiont acquisition in lucinid clams. *Environ. Microbiol.* 14, 1584–1595.

Herry, A., Diouris, M., and Le Pennec, M. (1989). Chemoautotrophic symbionts and translocation of fixed carbon from bacteria to host tissues in the littoral bivalve *Loripes lucinalis* (Lucinidae). *Mar. Biol.* 101, 305–312.

Hopsu-Havu, V.K., Arstila, A.U., Helminen, H.J., Kalimo, H.O., and Glenner, G.G. (1967). Improvements in the method for electron microscopic localization of aryl sulphatase activity. *Histochemie*.8, 54.

Johnson, M.A., and Fernandez, C. (2001). Bacterial symbiosis in *Loripes lucinalis* (Mollusca: Bivalvia) with comments on reproductive strategy. *J. Mar. Biol. Assoc. UK* 81, 251–257.

Jørgensen, B.B. (1982). Mineralization of organic matter in the sea bed - the role of sulphate reduction. *Nature* 296, 643–645.

Kádár, E., Davis, S.A., and Lobo-da-Cunha, A. (2007). Cytoenzymatic investigation of intracellular digestion in the symbiont-bearing hydrothermal bivalve *Bathymodiolus azoricus*. *Mar. Biol.* 153, 995–1004.

Lechaire, J.-P., Frébourg, G., Gaill, F., and Gros, O. (2008). In situ characterization of sulphur in gill-endosymbionts of the shallow water lucinid *Codakia orbicularis* (Linné, 1758) by high-pressure cryofixation and EFTEM microanalysis. *Mar. Biol.* 154, 693–700.

Liberge, M., Gros, O., and Frenkiel, L. (2001). Lysosomes and sulfide-oxidizing bodies in the bacteriocytes of *Lucina pectinata*, a cytochemical and microanalysis approach. *Mar. Biol.* 139, 401–409.

Löffler, B.-M., Hesse, B., and Kunze, H. (1984). A combined assay of three lysosomal marker enzymes: Acid phosphatase, β -d-glucuronidase and β -N-acetyl-d-hexosaminidase. *Anal. Biochem.* 142, 312–316.

Narbaiz, R. (1995). Cytochemical staining methods for electron microscopy, by P. R. Lewis and D. P. Knight. vol. 14 in “practical methods in electron microscopy,” A. M. glauert, ed., elsevier, Amsterdam, London, New york, Tokyo, 1992, 321 pp.

Ott, J., Rieger, R., and Enderes, F. (1982). New Mouth less Interstitial Worms from the Sulfide System: Symbiosis with Prokaryotes. *Mar. Ecol.* what is the volume number?? 313–333.

Patel, S., and Patel, B. (1985). Effect of environmental parameters on lysosomal marker enzymes in the tropical blood clam *Anadara granosa*. *Mar. Biol.* 85, 245–252.

Pipe, R.K., and Moore, M.N. (1985). The ultrastructural localization of lysosomal acid hydrolases in developing oocytes of the common marine mussel *Mytilus edulis*. *Histochem. J.* 17, 939–949.

Reid, R. (1990). Evolutionary implications of sulphide oxidizing symbioses in bivalves. In: The Bivalvia: Proceedings of a Memoriam Symposium in Honour of Sir C.M. Yonge, Edingburgh, 1986, B. Morton, ed, Hong Kong University Press 127-140.

Reynolds, L.K., Berg, P., and Zieman, J.C. (2007). Lucinid clam influence on the biogeochemistry of the seagrass *Thalassia testudinum* sediments. *Estuaries Coasts* 30, 482–490.

Rogers, D. (1960). Acid phosphatases of *E. coli*. *Arch. Biochem. Biophys.* 89, 97–104.

Schweimanns, M., and Felbeck, H. (1985). Significance of the occurrence of chemoautotrophic bacterial endosymbionts in lucinid clams from Bermuda. *Mar. Ecol.-Prog. Ser.* 24, 113–120.

Stewart, F.J., and Cavanaugh, C.M. (2006). Symbiosis of thioautotrophic bacteria with *Riftia pachyptila*. *Prog. Mol. Subcell. Biol.* 41, 197–225.

Streams, M.E., Fisher, C.R., and Fiala-Medioni, A. (1997). Methanotrophic symbiont location and fate of carbon incorporated from methane in a hydrocarbon seep mussel. *Mar. Biol.* 129, 465–476.

Taylor, J.D., and Glover, Emily A (1997). The lucinid bivalve genus *Cardiolucina* systematics anatomy and relationships. *Bull. Nat. Hist. Mus. Zool. Ser.* **what is the volume number??** 93–122.

Van Dover, C.L. (2000). The ecology of deep-sea hydrothermal vents (Princeton University Press).

Figure 1: Cytochemical detection of lysosomal enzymes arylsulfatase and acid phosphatase in thin gill sections of *Codakia orbiculata*.

A-B: Very few bacteriocytes from freshly collected individuals presented a couple of bacterial endosymbionts (*) partially degraded due to the activity of arylsulfatase (A) and to acid phosphatase (B). **C-D:** In individuals starved for two weeks, the number of bacteria degraded by lysosomal enzymes like acid phosphatase increases with numerous bacteriocytes affected. In each bacteriocyte (BC), several bacteria (b) are degraded by this enzyme suggesting a high lysosomal activity from host. In the same time, other gill cells (intercalary cell (IC) and granule cell (GC)) devoid of bacterial symbionts present no lysosomal activity. **E:** In bivalves starved for two month, most of the bacteria inside the bacteriocytes appear to be affected by arylsulfatase degradation. **F:** EDX spectrum obtained from cytochemical detection of acid phosphatase by lead precipitation from black areas located inside bacterial endosymbionts indicated by black arrows in pictures C and D. Copper originates from the grid bars supporting the thin section.

(BC: bacteriocytes; GC: granule cells; IC: intercalary cells; mv: microvilli; N: nucleus)

Figure 2: Cytochemical detection of lysosomal enzymes arylsulfatase and acid phosphatase in thin gill sections of *Codakia orbicularis*.

A-B: Conversely to *C. orbiculata*, lysosomal activity in *C. orbicularis* was weak or absent in freshly collected individuals and even after three weeks of starvation, no activity was detected either for arylsulfatase (A) or for acid phosphatase (B). **C-D:** Acid phosphatase activity from six weeks starved individuals concerns a large number of bacteria within most of the bacteriocytes composing each gill filament. In each bacteriocyte (BC), several bacteria (b) are degraded simultaneously by this enzyme confirming a strong lysosomal activity from the host cell. In the same time, granule cell (GC), which are normally devoid of bacterial endosymbionts, present no lysosomal activity. **E:** In bivalves starved for six weeks, all the symbionts appear to be affected by arylsulfatase degradation as shown by the numerous black areas observed through the cytoplasm of the bacteriocytes. **F:** EDX spectrum obtained from cytochemical detection of arylsulfatase activity by barium precipitation from black areas

(black arrows) in pictures C-E. Copper originates from the grid bars supporting the thin section.

(BC: bacteriocytes; GC: granule cells; IC: intercalary cells; mv: microvilli; N: nucleus)