

ON THE CONTROL OF ILL-POSED DISTRIBUTED PARAMETER SYSTEMS

René Dorville, O Nakoulima, A Omrane

► **To cite this version:**

René Dorville, O Nakoulima, A Omrane. ON THE CONTROL OF ILL-POSED DISTRIBUTED
PARAMETER SYSTEMS. ESAIM: Proceedings, EDP Sciences, 2007, 17, pp.50 - 66. hal-01683792

HAL Id: hal-01683792

<https://hal.univ-antilles.fr/hal-01683792>

Submitted on 14 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE CONTROL OF ILL-POSED DISTRIBUTED PARAMETER SYSTEMS

R. DORVILLE¹, O. NAKOULIMA² AND A. OMRANE³

Abstract. We show that the so-called low-regret (or least-regret) control by J. L. Lions [8] fits on the control of ill-posed problems. At each time, we give the characterization of the so-called no-regret control by means of singular optimality systems. For the backward heat ill-posed problem, no Slater hypothesis is assumed on the admissible set of controls \mathcal{U}_{ad} .

This work is two pieces, and two methods are considered : the regularization method and the null-controllability method. For the first method, a zero order corrector is used, while for the second method, the passage to the limit is easy. The results presented here generalize the works in [2, 3] to the no-regret control.

Résumé. On montre que le contrôle à moindres regrets de J. L. Lions [8] est bien adapté pour le contrôle des problèmes mal posés. A chaque fois, on donne une caractérisation du contrôle sans regret par le moyen de systèmes d'optimalité singuliers. Pour le problème de chaleur rétrograde mal posé, aucune hypothèse de type Slater sur l'ensemble des contrôles admissibles \mathcal{U}_{ad} n'est nécessaire.

Ce travail est divisé en deux parties, et deux méthodes sont considérées: la méthode de régularisation et celle de la contrôlabilité à zéro. Pour la première méthode, un correcteur d'ordre zéro est utilisé, alors que pour la seconde méthode le passage à la limite est simple. Les résultats présentés ici généralisent les travaux dans [2, 3] au contrôle sans regret.

CONTENTS

1. Introduction	51
1.1. Preliminaries	51
1.2. Existence of a solution for the ill-posed heat problem	52
1.3. The optimal control question	53
2. The regularization approach	54
2.1. A priori estimates. The low-regret control	57
2.2. Zero-order corrector	59
2.3. Passage to the limit	61
3. The controllability approach	62
3.1. The low-regret control for the null-controllable problem	63
3.2. Passage to the limit	65
References	66

¹ Université Antilles-Guyane, Laboratoire d'Analyse, Optimisation et Contrôle (AOC), UFR Sciences Exactes et Naturelles, Campus de Fouillole, 97159 Pointe à Pitre. E-mail: rene.dor@wanadoo.fr

² onakouli@univ-ag.fr

³ aomrane@univ-ag.fr

1. INTRODUCTION

The solutions of singular optimal control problems, particularly of ill-posed problems are not regular. As a consequence, the characterization of such solutions has not been deeply investigated. Moreover, the assumptions on the control problem are frequently chosen in such a way that some standard methods can be applied to derive the existence of solutions.

We consider here the prototype of ill-posed problems : the ill-posed backward heat problem, with controls v in a non-empty closed convex subset \mathcal{U}_{ad} of the Hilbert space L^2 , and a cost functional.

It is well known that if we add the Slater type hypothesis as : \mathcal{U}_{ad} has a non-empty interior, then we ensure the existence of an optimal control, using standard methods (see Lions [6]). But, we do not know if this hypothesis is unavoidable !

One way to deal with this problem -instead of answering the question-, is to propose another approach, where the Slater hypothesis on \mathcal{U}_{ad} is not needed. Convex cones as $(L^2)^+$ which are empty interiored may be used as set of feasible controls v . We here use the regularization approach in a first part, and the null-controllability approach in a second part. In both methods, a new datum is introduced. These data are supposed to be chosen as largely as possible, say in a vector space. That is the data are *incomplete*, and in this case, the standard methods as the penalization method are not adapted (see Lions [8], and Nakoulima-Omrane-Velin [13] [14]). We then seek for the low-regret control of the distributed system of incomplete data obtained. And thus, the no-regret control of the original problem appears naturally by taking some limit, which is possible without the Slater hypothesis.

Roughly speaking, the low-regret control u^γ satisfies to the following inequality :

$$J(u^\gamma, g) \leq J(0, g) + \gamma \|g\|_Y^2 \quad \forall g \text{ in a Hilbert space } Y,$$

where γ is a small positive parameter (g being the pollution or incomplete data). With the low-regret control we admit the possibility of making a choice of controls v 'slightly worse' than by doing better than $v = 0$ -but 'not much' if we choose γ small enough- compared to the worst things that could happen with the 'pollution' g . In the no-regret concept, we search for the control u , if it exists, which makes things better than $v = 0$, for any given perturbation parameter. It is the limit when $\gamma \rightarrow 0$, of the family of low-regret controls u^γ .

This concept is previously introduced by Savage [17] in statistics. Lions was the first to apply it to control distributed systems of incomplete data, motivated by a number of applications in economics, and ecology as well (see for instance [9], [10]).

In [13] (see also [14]), Nakoulima *et al.* give a precise optimality system (which is a singular optimality system). In [14], the no-regret control for problems of incomplete data, in both the stationary and evolution cases is characterized. A number of applications is given too.

In the litterature mentioned above, the only regular problems are considered. Moreover, the set of controls was a Hilbert space. In this article, we generalize the study to the control of ill-posed problems, where the controls are in a closed convex subset of a Hilbert space only. Without loss of generality, we consider the typical ill-posed problems: *the ill-posed backward heat problem*.

1.1. Preliminaries

Consider an open domain $\Omega \subset \mathbb{R}^N$ with smooth boundary $\partial\Omega$, and denote by $Q = (0, T) \times \Omega$, and by $\Sigma = (0, T) \times \partial\Omega$. Then it is well known that the following heat system :

$$z' - \Delta z = v \quad \text{in } Q, \tag{1}$$

$$z = 0 \quad \text{on } \Sigma, \tag{2}$$

$$z(0) = 0 \quad \text{in } \Omega, \tag{3}$$

is well-posed. Here, $z = z(t, x)$ is the state solution and $v = v(t, x) \in L^2(Q)$. Moreover, any solution z of (1)-(3) is a.e. equal to a continuous function from $[0, T]$ to $H^{-1}(\Omega)$ and we have

$$z' \in L^2(]0, T[; H^{-2}(\Omega)) \quad \text{and} \quad z|_{\Sigma} \in H^{-1}\left(]0, T[; H^{-\frac{1}{2}}(\partial\Omega)\right).$$

But, the above system does not admit a solution for arbitrary data (3). Indeed, in the case of final data, i.e. replacing (3) by

$$z(T) = 0 \quad \text{in } \Omega, \quad (4)$$

there is no solution for the *backward* heat problem, even for regular control v , as we can see in the following one-dimensional example.

A counter-example - For $\Omega =]0, \pi[$, $T = 1$, consider the backward heat system

$$\begin{cases} \frac{\partial z}{\partial t} - \frac{\partial^2 z}{\partial x^2} = v & \text{in }]0, \pi[\times]0, 1[, \\ z(0, t) = z(\pi, t) = 0 & \text{in }]0, 1[, \\ z(x, 1) = 0 & \text{in }]0, \pi[, \end{cases} \quad (5)$$

where $v \in L^2(]0, 1[; L^2(]0, \pi[))$ is the uniformly convergent series

$$v(t, x) = \sqrt{\frac{2}{\pi}} \sum_{m \geq 1} \frac{\sin mx}{m^2}.$$

If $z \in L^2(]0, 1[; L^2(]0, \pi[))$ is a solution to (5), then $z(t, x) = \sum_{m \geq 1} z_m(t) w_m(x)$, where $w_m(x) = \sqrt{\frac{2}{\pi}} \sin mx$ (w_m is an eigenfunction for $-\frac{\partial^2}{\partial x^2}$ related to the eigenvalue m^2), then we have

$$\begin{cases} \frac{dz_m}{dt}(t) + m^2 z_m(t) = \frac{1}{m^2} & \text{in }]0, 1[, \\ z_m(1) = 0, \end{cases}$$

so that

$$z_m(t) = \frac{1}{m^2} \int_1^t e^{m^2(s-t)} ds = \frac{1}{m^4} \left(1 - e^{m^2(1-t)}\right).$$

For every $t \in [0, 1[$, we then obtain $\|z\|_{L^2(]0, \pi[)}^2 = \sum_{m \geq 1} \left| \frac{1}{m^4} \left(1 - e^{m^2(1-t)}\right) \right|^2$. But,

$$\lim_{m \rightarrow +\infty} \left| \frac{1}{m^4} \left(1 - e^{m^2(1-t)}\right) \right|^2 = +\infty, \quad \forall t \in [0, 1[.$$

Hence, the series diverges and the solution z of (5) does not exist.

1.2. Existence of a solution for the ill-posed heat problem

We consider the following backward heat problem :

$$\begin{cases} z' - \Delta z = v & \text{in } Q, \\ z = 0 & \text{on } \Sigma, \\ z(T) = 0 & \text{in } \Omega, \end{cases} \quad (6)$$

which is a prototype of ill-posed problems.

Remark 1.1. The backward problem (6) has a unique solution z but for v in a some dense subset of $L^2(Q)$.

Indeed, for example, we can consider the vector space

$$V = \left\{ w = \sum_{i=1}^N \lambda_i w_i : -\Delta w_i = 0, \quad w_i = 0 \text{ on } \partial\Omega, \quad \text{and} \quad w_i \in L^2(\Omega) \right\}, \quad (7)$$

there is $f \in L^2(]0, T[)$ and $w \in V$ such that

$$v(t, x) = f(t) \left(\sum_{i=1}^N \lambda_i w_i(x) \right),$$

for given $v \in L^2(]0, T[) \otimes V$ (which is dense in $L^2(Q)$). It suffices to take z of the form $z(t, x) = \zeta(t)w(x)$, $\zeta = (\zeta_1, \dots, \zeta_N)$. So, ζ_i is solution of

$$\begin{cases} \frac{\partial \zeta_i}{\partial t} - \lambda \zeta_i = f & \text{in }]0, T[, \\ \zeta_i(0) = 0 & \text{in } \Omega, \end{cases}$$

which defines ζ in a unique manner.

1.3. The optimal control question

Consider $v \in \mathcal{U}_{ad}$, \mathcal{U}_{ad} a non-empty closed convex subset of the Hilbert space of controls $L^2(Q)$, and the quadratic function

$$J(v, z) = \left\| z - z_d \right\|_{L^2(Q)}^2 + N \left\| v \right\|_{L^2(Q)}^2, \quad (8)$$

where $z_d \in L^2(Q)$, $N > 0$, and where $\left\| \cdot \right\|_X$ is the norm on the corresponding Hilbert space X .

If a pair $(v, z) \in \mathcal{U}_{ad} \times L^2(Q)$ satisfying (6) exists, then it is called a control-state feasible pair. Denote by \mathcal{X}_{ad} the set of admissible pairs for which (6) holds. We suppose in what follows that \mathcal{X}_{ad} is non-empty. Then for every $(v, z) \in \mathcal{X}_{ad}$, we associate the *cost function* defined by (8), and we consider the *optimal control problem* :

$$\inf J(v, z), \quad (v, z) \in \mathcal{U}_{ad} \times L^2(Q) \quad (9)$$

which has a unique solution (u, y) that we should characterize.

Lemma 1.2. *The problem (9) has one only solution (u, y) called the optimal pair.*

Proof. The functional $J : L^2(Q) \times L^2(Q) \rightarrow \mathbb{R}$ is a lower semi-continuous function, strictly convex, and coercitive. Hence there is a unique admissible pair (u, y) solution to (9). A classical method to control the system (6) and (8) is the well-known penalization method, which consists in approximating (u, y) by the solution of some penalized problem. More precisely, for $\varepsilon > 0$ we define the penalized cost function

$$J_\varepsilon(v, z) = J(v, z) + \frac{1}{2\varepsilon} \|z' - \Delta z - v\|_{L^2(Q)}^2.$$

The optimal pair $(u_\varepsilon, y_\varepsilon)$ then converges to (u, y) when $\varepsilon \rightarrow 0$.

The optimality conditions of Euler-Lagrange for $(u_\varepsilon, y_\varepsilon)$ are the following :

$$\frac{d}{dt} J_\varepsilon(u_\varepsilon, y_\varepsilon + t(z - y_\varepsilon))|_{t=0} = 0, \quad \forall z \in \mathcal{F} \quad (10)$$

and

$$\frac{d}{dt} J_\varepsilon(u_\varepsilon + t(v - u_\varepsilon), y_\varepsilon)|_{t=0} \geq 0 \quad \forall v \in \mathcal{U}_{ad}, \quad (11)$$

then an optimality system is obtained by the introduction of the *adjoint state*

$$p_\varepsilon = -\frac{1}{\varepsilon}(y_\varepsilon' - \Delta y_\varepsilon - u_\varepsilon).$$

A priori estimates (consisting in bounding p_ε in $L^2(Q)$) have to be obtained, which allows the passage to the limit under some hypothesis : For the problem (6) and (8)-(9), J. L. Lions obtained in [6] a singular optimality system, under the extra hypothesis of Slater type :

$$\mathcal{U}_{ad} \text{ has a non-empty interior.} \quad (12)$$

More precisely, we have the following theorem due to Lions :

Theorem 1.3. (J.L. Lions [6]) *Under hypothesis (12), there is a unique $(u, y, p) \in \mathcal{U}_{ad} \times L^2(Q) \times L^2(Q)$, solution to the optimal control problem (6) and (8)-(9). Moreover, this solution is characterized by the following singular optimality system (SOS) :*

$$\begin{cases} y' - \Delta y = u, & -p' - \Delta p = y - z_d & \text{in } Q, \\ y(T) = 0, & p(0) = 0 & \text{in } \Omega, \\ y = 0, & p = 0 & \text{on } \Sigma \end{cases} \quad (13)$$

with the variational inequality

$$(p + Nu, v - u)_{L^2(Q)} \geq 0 \quad \forall v \in \mathcal{U}_{ad}. \quad (14)$$

Remark 1.4. Of course when $\mathcal{U}_{ad} = L^2(Q)$, the hypothesis (12) is satisfied and then the above theorem holds. In some applications, the Slater hypothesis (12) is not satisfied, as when $\mathcal{U}_{ad} = (L^2(Q))^+$ which has an empty interior. In the following, we propose alternative approach which does not assume (12).

2. THE REGULARIZATION APPROACH

In this section, some elliptic regularization of the ill-posed parabolic problem (6) is proposed. We obtain some well-posed problem but with a new *unknown* datum. We then let the classical control notion away, to consider the one of no-regret control by Lions [8], and recently developed by Nakoulima *et al.* in [13] [14].

For any $\varepsilon > 0$, we consider the regularized problem :

$$\begin{cases} -\varepsilon z_\varepsilon'' + z_\varepsilon' - \Delta z_\varepsilon = v & \text{in } Q, \\ z_\varepsilon(0) = g & \text{in } \Omega, \\ z_\varepsilon(T) = 0 & \text{in } \Omega, \\ z_\varepsilon = 0 & \text{on } \Sigma, \end{cases} \quad (15)$$

where $g \in L^2(\Omega)$.

It is clear that for any $\varepsilon > 0$, and any data (v, g) , there is a unique state solution $z_\varepsilon = z_\varepsilon(v, g)$ of the elliptic problem (15) (see Lions-Magenes [12] for example), for which we associate the cost given by :

$$J_\varepsilon(v, g) = \left\| z_\varepsilon(v, g) - z_d \right\|_{L^2(Q)}^2 + N \left\| v \right\|_{L^2(Q)}^2, \quad g \in L^2(\Omega). \quad (16)$$

We are concerned with the optimal control of the problem (15)–(16). Clearly we want

$$\inf_{v \in \mathcal{U}_{ad}} J_\varepsilon(v, g) \quad \forall g \in L^2(\Omega). \quad (17)$$

If $g \in G \subset L^2(\Omega)$ with G finite, then we can give a Pareto sense to (17). But if G is not finite, then the above minimization problem has no sense. One natural idea is to consider the following minimization problem :

$$\inf_{v \in \mathcal{U}_{ad}} \left(\sup_{g \in L^2(\Omega)} J_\varepsilon(v, g) \right),$$

but from above, we see that J_ε is not bounded, since $\sup_{g \in L^2(\Omega)} J_\varepsilon(v, g) = +\infty$. The idea of Lions is then to look for controls v -if they exist- such that

$$J_\varepsilon(v, g) \leq J_\varepsilon(0, g) \quad \forall g \in L^2(\Omega),$$

and thus

$$J_\varepsilon(v, g) - J_\varepsilon(0, g) \leq 0 \quad \forall g \in L^2(\Omega).$$

Those controls *doing better than* $v = 0$ for every pollution g are called no-regret controls.

Definition 2.1. We say that $u \in \mathcal{U}_{ad}$ is a no-regret control for (15)–(16) if u is a solution to the following problem:

$$\inf_{v \in \mathcal{U}_{ad}} \left(\sup_{g \in L^2(\Omega)} (J_\varepsilon(v, g) - J_\varepsilon(0, g)) \right). \quad (18)$$

Lemma 2.2. For any $v \in \mathcal{U}_{ad}$ we have

$$J_\varepsilon(v, g) - J_\varepsilon(0, g) = J_\varepsilon(v, 0) - J_\varepsilon(0, 0) + 2\langle \xi_\varepsilon'(0), g \rangle_{L^2(\Omega)} \quad \forall g \in L^2(\Omega), \quad (19)$$

where ξ_ε satisfies to :

$$-\xi_\varepsilon' - \varepsilon \xi_\varepsilon'' - \Delta \xi_\varepsilon = y_\varepsilon(v, 0) \text{ in } Q, \quad \xi_\varepsilon(0) = \xi_\varepsilon(T) = 0 \text{ in } \Omega, \quad \xi_\varepsilon = 0 \text{ on } \Sigma. \quad (20)$$

Proof. A direct computation gives

$$J_\varepsilon(v, g) - J_\varepsilon(0, g) = J_\varepsilon(v, 0) - J_\varepsilon(0, 0) + 2\langle z_\varepsilon(v, 0); z_\varepsilon(0, g) \rangle_{L^2(Q)}.$$

Following the the Green formula we get

$$\langle z_\varepsilon(v, 0); z_\varepsilon(0, g) \rangle = \varepsilon \langle \xi_\varepsilon'(0); g \rangle_{L^2(Q)},$$

where ξ_ε is given by (20).

Remark 2.3. Of course the problem (18) is defined only for the controls $v \in \mathcal{U}_{ad}$ such that

$$\sup_{g \in L^2(\Omega)} (J_\varepsilon(v, g) - J_\varepsilon(0, g)) < \infty.$$

From (19) this is achieved iff $v \in K$, where $K = \{w \in \mathcal{U}_{ad}, \langle \xi_\varepsilon(w), g \rangle = 0 \forall g \in L^2(\Omega)\}$. This set is hard to characterize. As in [13], for any $\gamma > 0$, we define the control in the low-regret sense.

Definition 2.4. The low-regret control for (15)–(16), is the solution to the following perturbed system :

$$\inf_{v \in \mathcal{U}_{\text{ad}}} \left(\sup_{g \in L^2(\Omega)} \left(J_\varepsilon(v, g) - J_\varepsilon(0, g) - \gamma \|g\|_{L^2(\Omega)}^2 \right) \right). \quad (21)$$

Remark that :

$$\begin{aligned} & \inf_{v \in \mathcal{U}_{\text{ad}}} \left(\sup_{g \in L^2(\Omega)} \left(J_\varepsilon(v, g) - J_\varepsilon(0, g) - \gamma \|g\|_{L^2(\Omega)}^2 \right) \right) \\ &= \inf_{v \in \mathcal{U}_{\text{ad}}} \left(J_\varepsilon(v, 0) - J_\varepsilon(0, 0) + \left(\sup_{g \in L^2(\Omega)} 2 \langle z_\varepsilon(v, 0); z_\varepsilon(0, g) \rangle_{L^2(Q)} - \gamma \|g\|_{L^2(\Omega)}^2 \right) \right). \end{aligned}$$

Thanks to Legendre transform, we get the classical control problem :

$$\inf_{v \in \mathcal{U}_{\text{ad}}} \mathcal{J}_\varepsilon^\gamma(v) \quad (22)$$

where

$$\mathcal{J}_\varepsilon^\gamma(v) = J_\varepsilon(v, 0) - J_\varepsilon(0, 0) + \frac{\varepsilon^2}{\gamma} \left\| \xi'_\varepsilon(T, v) \right\|_{L^2(\Omega)}^2, \quad (23)$$

and where ξ_ε satisfies (20).

Remark 2.5. As we can see here, the low-regret control method allows to transform systematically problems with uncertainty into standard control problems. We can then use the Euler–Lagrange method.

We can replace now (21) by (22) and (23) for the low-regret control.

Lemma 2.6. *The problem (15) with control (22)–(23) has one only solution u_ε^γ , called the ‘approximate’ low-regret control.*

Proof. We have $\mathcal{J}_\varepsilon^\gamma(v) \geq -J_\varepsilon(0, 0) = -\|z_d\|_{L^2(\Omega)}^2 \forall v \in \mathcal{U}_{\text{ad}}$. Then $d = \inf_{v \in \mathcal{U}_{\text{ad}}} \mathcal{J}_\varepsilon^\gamma(v) > -\infty$. Let v_n be some minimizing sequence (such that $d = \lim_{n \rightarrow \infty} \mathcal{J}_\varepsilon^\gamma(v_n)$). We have

$$-\|z_d\|_{L^2(\Omega)}^2 \leq \mathcal{J}_\varepsilon^\gamma(v_n) = J_\varepsilon(v_n, 0) - J_\varepsilon(0, 0) + \frac{1}{\gamma} \left\| \xi'_\varepsilon(0) \right\|_{L^2(\Omega)}^2 \leq d_\gamma + 1.$$

Then we deduce the bounds

$$\|v_n\|_{L^2(Q)} \leq c, \quad \frac{1}{\sqrt{\gamma}} \left\| \xi'_\varepsilon(v_n)(0) \right\|_{L^2(\Omega)} \leq c, \quad \left\| y_\varepsilon(v_n, 0) - z_d \right\|_{L^2(Q)} \leq c,$$

where the constant c is independent of n .

There exists $u_\varepsilon^\gamma \in \mathcal{U}_{\text{ad}}$ such that $v_n \rightharpoonup u_\varepsilon^\gamma$ weakly in \mathcal{U}_{ad} (which is closed). Also, $y_\varepsilon(v_n, 0) \rightarrow y_\varepsilon(u_\varepsilon^\gamma, 0)$ (continuity w.r.t the data). Now since $\mathcal{J}_\varepsilon^\gamma$ is strictly convex, u_ε^γ is unique.

Proposition 2.7. *The 'approximate' low-regret control u_ε^γ is characterized by the unique quadruplet $\{u_\varepsilon^\gamma, y_\varepsilon^\gamma, \rho_\varepsilon^\gamma, p_\varepsilon^\gamma\}$, solution to the system :*

$$\begin{cases} y_\varepsilon^{\gamma'} - \varepsilon y_\varepsilon^{\gamma''} - \Delta y_\varepsilon^\gamma = u_\varepsilon^\gamma, & \rho_\varepsilon^{\gamma'} - \varepsilon \rho_\varepsilon^{\gamma''} - \Delta \rho_\varepsilon^\gamma = 0, \text{ and} \\ -p_\varepsilon^{\gamma'} - \varepsilon p_\varepsilon^{\gamma''} - \Delta p_\varepsilon^\gamma = y_\varepsilon^\gamma - z_d + \rho_\varepsilon^\gamma & \text{in } Q, \\ y_\varepsilon^\gamma(0) = y_\varepsilon^\gamma(T) = 0, & \rho_\varepsilon^\gamma(T) = 0, & \rho_\varepsilon^\gamma(0) = \frac{\varepsilon}{\gamma} \xi_\varepsilon^{\gamma'}(0), \\ p_\varepsilon^\gamma(0) = p_\varepsilon^\gamma(T) = 0 & \text{in } \Omega, \\ y_\varepsilon^\gamma = 0, & \rho_\varepsilon^\gamma = 0, & p_\varepsilon^\gamma = 0 \text{ on } \Sigma, \end{cases}$$

with (20),

and the variational inequality :

$$\langle p_\varepsilon^\gamma + N u_\varepsilon^\gamma, v - u_\varepsilon^\gamma \rangle \geq 0, \quad \forall v \in \mathcal{U}_{ad}.$$

Proof. The first order Euler condition for (22) and (23) gives :

$$\langle y_\varepsilon^\gamma - z_d, y_\varepsilon(w, 0) \rangle_{L^2(Q) \times L^2(Q)} + N \langle u_\varepsilon^\gamma, w \rangle_{L^2(Q) \times L^2(Q)} + \langle \frac{\varepsilon^2}{\gamma} \xi_\varepsilon^{\gamma'}(0), \xi_\varepsilon'(0, w) \rangle_{L^2(\Omega) \times L^2(\Omega)} \geq 0,$$

where $y_\varepsilon^\gamma = y_\varepsilon(u_\varepsilon^\gamma, 0)$, and $\xi_\varepsilon^\gamma = \xi_\varepsilon(u_\varepsilon^\gamma, 0)$. We then define $\rho_\varepsilon^\gamma = \rho_\varepsilon(u_\varepsilon^\gamma, 0)$ which is the solution of $\rho_\varepsilon^{\gamma'} - \varepsilon \rho_\varepsilon^{\gamma''} - \Delta \rho_\varepsilon^\gamma = 0$, $\rho_\varepsilon^\gamma(0) = (\varepsilon/\gamma) \xi_\varepsilon^{\gamma'}(0)$, $\rho_\varepsilon^\gamma(T) = 0$, and $\rho_\varepsilon^\gamma = 0$ on Σ , such that :

$$\langle \frac{\varepsilon^2}{\gamma} \xi_\varepsilon^{\gamma'}(0), \xi_\varepsilon'(0, w) \rangle_{L^2(\Omega) \times L^2(\Omega)} = \langle \varepsilon \rho_\varepsilon^\gamma(0), \xi_\varepsilon'(0, w) \rangle_{L^2(\Omega) \times L^2(\Omega)} = \langle \rho_\varepsilon^\gamma, y_\varepsilon(w, 0) \rangle_{L^2(Q) \times L^2(Q)}$$

by the Green formula.

Introduce now the adjoint state $p_\varepsilon^\gamma = p_\varepsilon(u_\varepsilon^\gamma, 0)$ as follows : we solve $-p_\varepsilon^{\gamma'} - \varepsilon p_\varepsilon^{\gamma''} - \Delta p_\varepsilon^\gamma = y_\varepsilon^\gamma - z_d + \rho_\varepsilon^\gamma$, $p_\varepsilon^\gamma(T) = p_\varepsilon^\gamma(0) = 0$, and $p_\varepsilon^\gamma = 0$ on Σ . Hence we have

$$\langle y_\varepsilon^\gamma - z_d + \rho_\varepsilon^\gamma, y_\varepsilon(w, 0) \rangle_{L^2(Q) \times L^2(Q)} = \langle p_\varepsilon^\gamma, w \rangle_{L^2(Q) \times L^2(Q)}.$$

Finally,

$$\langle p_\varepsilon^\gamma + N u_\varepsilon^\gamma, w \rangle_{L^2(Q) \times L^2(Q)} \geq 0.$$

2.1. A priori estimates. The low-regret control

In this section we give the S.O.S for the low-regret control of the backward heat equation. Estimates are given in the following statement :

Proposition 2.8. *There is some positive constant C , and, for any small $\eta > 0$, there is some constant $C_\eta > 0$ such that :*

$$\|u_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \|y_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \frac{\varepsilon}{\sqrt{\gamma}} \|\xi_\varepsilon^{\gamma'}(0)\|_{L^2(\Omega)} \leq C, \quad (24)$$

$$\varepsilon \|y_\varepsilon^{\gamma'}\|_{L^2(Q)} + \|y_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \|y_\varepsilon^{\gamma'}\|_{L^2([0, T-\eta]; H^{-1}(\Omega))} \leq C_\eta, \quad (25)$$

and,

$$\varepsilon \|\xi_\varepsilon^{\gamma'}\|_{L^2(Q)} + \|\xi_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \|\xi_\varepsilon^{\gamma'}\|_{L^2([0,T];H^{-1}(\Omega))} \leq C_\eta, \quad (26)$$

$$\varepsilon \|\rho_\varepsilon^{\gamma'}\|_{L^2(Q)} + \|\rho_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \|\rho_\varepsilon^{\gamma'}\|_{L^2([0,T-\eta];H^{-1}(\Omega))} \leq C_\eta, \quad (27)$$

$$\varepsilon \|p_\varepsilon^{\gamma'}\|_{L^2(Q)} + \|p_\varepsilon^\gamma\|_{L^2(Q)} \leq C, \quad \|p_\varepsilon^{\gamma'}\|_{L^2([0,T];H^{-1}(\Omega))} \leq C_\eta. \quad (28)$$

Proof. Since u_ε^γ is the approximate low-regret control

$$\mathcal{J}_\varepsilon^\gamma(u_\varepsilon^\gamma) \leq \mathcal{J}_\varepsilon^\gamma(v) \quad \forall v \in \mathcal{U}_{ad},$$

in the particular case where $v = 0$,

$$J_\varepsilon(u_\varepsilon^\gamma, 0) - J_\varepsilon(0, 0) + \frac{\varepsilon^2}{\gamma} \left\| \xi_\varepsilon'(u_\varepsilon^\gamma)(0) \right\|_{L^2(\Omega)}^2 \leq \frac{\varepsilon^2}{\gamma} \left\| \xi_\varepsilon'(u_\varepsilon^\gamma)(0) \right\|_{L^2(\Omega)}^2.$$

On the other hand $y_\varepsilon(0, 0)(t, x) = \xi_\varepsilon(0)(t, x) = 0$ in $[0, T] \times \bar{\Omega}$, then :

$$\left\| y_\varepsilon(u_\varepsilon^\gamma, 0) - z_d \right\|_{L^2(Q)}^2 + N \left\| u_\varepsilon^\gamma \right\|_{L^2(Q)}^2 + \left\| \frac{\varepsilon}{\sqrt{\gamma}} \xi_\varepsilon'(u_\varepsilon^\gamma)(0) \right\|_{L^2(\Omega)}^2 \leq \left\| z_d \right\|_{L^2(Q)}^2 = \text{constant}, \quad (29)$$

so (24) holds.

From the Poincaré formula, there is some constant $C_1 > 0$ such that :

$$\left(\varepsilon \left\| \frac{\partial y_\varepsilon^\gamma}{\partial t} \right\|_{L^2(Q)} + \|y_\varepsilon^\gamma\|_{L^2(Q)} \right) \|y_\varepsilon^\gamma\|_{L^2(Q)} \leq C_1 \|u_\varepsilon^\gamma\|_{L^2(Q)} \|y_\varepsilon^\gamma\|_{L^2(Q)},$$

thus, there is some constant $C_2 > 0$ such that

$$\varepsilon \left\| \frac{\partial y_\varepsilon^\gamma}{\partial t} \right\|_{L^2(Q)} + \|y_\varepsilon^\gamma\|_{L^2(Q)} \leq C_2,$$

that is the first part of (25). Now, we consider the equation $-\varepsilon \frac{\partial^2 y_\varepsilon}{\partial t^2} + \frac{\partial y_\varepsilon}{\partial t} = v + \Delta y_\varepsilon$, and denote by $g_\varepsilon = v + \Delta y_\varepsilon$.

Then g_ε stays in some bounded subset of $L^2(Q)$.

Consider some function $\varphi \in C^1([0, T])$ such that $\varphi(0) = 1$ and $\varphi(T) = 0$. Multiply both members of

$$-\varepsilon \frac{\partial^2 y_\varepsilon}{\partial t^2} + \frac{\partial y_\varepsilon}{\partial t} = g_\varepsilon$$

by $\varphi \frac{\partial y_\varepsilon}{\partial t}$ and integrate over the whole set Q . We have

$$-\frac{\varepsilon}{2} \int_0^T \varphi \frac{d}{dt} \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \varphi \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt = \int_0^T \varphi \left(g_\varepsilon, \frac{\partial y_\varepsilon}{\partial t} \right)_{L^2(\Omega)} dt. \quad (30)$$

And,

$$\int_0^T \varphi \frac{d}{dt} \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt = - \left\| \frac{\partial y_\varepsilon}{\partial t}(0) \right\|_{L^2(\Omega)}^2 - \int_0^T \frac{d\varphi}{dt} \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt,$$

thus

$$\frac{\varepsilon}{2} \left\| \frac{\partial y_\varepsilon}{\partial t}(0) \right\|_{L^2(\Omega)}^2 + \frac{\varepsilon}{2} \int_0^T \frac{d\varphi}{dt} \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt + \int_0^T \varphi \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt = \int_0^T \varphi \left(g_\varepsilon, \frac{\partial y_\varepsilon}{\partial t} \right)_{L^2(\Omega)} dt.$$

The second term is $O(1)$, so

$$\int_0^T \varphi \left\| \frac{\partial y_\varepsilon}{\partial t} \right\|_{L^2(\Omega)}^2 dt = \int_0^T \varphi \left(g_\varepsilon, \frac{\partial y_\varepsilon}{\partial t} \right)_{L^2(\Omega)} dt + O(1).$$

We finally deduce (25) from the triangular inequality.

The estimates (26), (27) and (28) follow easily.

Theorem 2.9. *The low-regret control $u^\gamma = \lim_{\varepsilon \rightarrow 0} u_\varepsilon^\gamma$ for the backwards heat equation (6) is characterized by the unique $\{u^\gamma, y^\gamma, \xi^\gamma, \rho^\gamma, p^\gamma\}$, solution to the system :*

$$\left\{ \begin{array}{l} y^{\gamma'} - \Delta y^\gamma = u^\gamma, \quad -\xi^{\gamma'} - \Delta \xi^\gamma = y^\gamma, \quad \rho^{\gamma'} - \Delta \rho^\gamma = 0, \quad \text{and} \\ -p^{\gamma'} - \Delta p^\gamma = y^\gamma - z_d + \rho^\gamma \quad \text{in } Q, \\ \\ y^\gamma(0) = 0, \quad \xi^\gamma(T) = 0, \quad \text{and} \\ \rho^\gamma(0) = \lambda^\gamma(0), \quad p^\gamma(T) = 0 \quad \text{in } \Omega, \\ \\ y^\gamma = 0, \quad \xi^\gamma = 0, \quad \rho^\gamma = 0, \quad p^\gamma = 0 \quad \text{on } \Sigma, \end{array} \right.$$

with the following weak limits

$$y^\gamma = \lim_{\varepsilon \rightarrow 0} y_\varepsilon^\gamma, \quad \xi^\gamma = \lim_{\varepsilon \rightarrow 0} \xi_\varepsilon^\gamma, \quad \rho^\gamma = \lim_{\varepsilon \rightarrow 0} \rho_\varepsilon^\gamma, \quad p^\gamma = \lim_{\varepsilon \rightarrow 0} p_\varepsilon^\gamma,$$

and the variational inequality :

$$\langle p^\gamma + Nu^\gamma, v - u^\gamma \rangle \geq 0 \quad \forall v \in \mathcal{U}_{ad},$$

where

$$u^\gamma, y^\gamma, p^\gamma, \rho^\gamma, \xi^\gamma \in L^2(]0, T[; L^2(\Omega)), \quad \lambda^\gamma(0) \in L^2(\Omega).$$

Proof. We use the estimates of proposition 2.8. From (29), we deduce the following limits :

$$\begin{array}{lll} u_\varepsilon^\gamma & \rightharpoonup & u^\gamma \quad \text{weakly in } \mathcal{U}_{ad}, \\ y_\varepsilon^\gamma & \rightharpoonup & y^\gamma \quad \text{weakly in } L^2(Q), \\ \frac{\varepsilon}{\sqrt{\gamma}} \frac{\partial \xi_\varepsilon(u_\varepsilon^\gamma)}{\partial t}(0) & \rightharpoonup & \lambda^\gamma(0) \quad \text{weakly in } L^2(\Omega), \end{array} \quad (31)$$

(up to extract a subsequences (u_ε^γ) , (y_ε^γ) and $\left(\frac{\varepsilon}{\sqrt{\gamma}} \frac{\partial \xi_\varepsilon(u_\varepsilon^\gamma)}{\partial t}(T)\right)$).

For fixed $\gamma > 0$, the adjoint state p_ε^γ depends boundedly on ε (from (28)).

2.2. Zero-order corrector

For the no-regret optimal control to the original problem, we now introduce the notion of corrector of order 0 of Lions [7] for elliptic regularizations, instead of using the Slater hypothesis (12). Indeed, as it is well known, the passage to the limit gives no information on $y^\gamma(T)$. Moreover, $y^\gamma(T) \neq 0$ in general (see Dorville [1] for details).

We assume the following :

$$y^\gamma(T) \in H_0^1(\Omega), \quad y^{\gamma'} \in L^2(Q). \quad (32)$$

Denote by

$$\mathcal{V} = \{\varphi \in L^2(]0, T[; H_0^1(\Omega)) \text{ such that } \varphi' \in L^2(Q)\},$$

and by

$$\mathcal{V}_0 = \{\varphi \in \mathcal{V} \text{ such that } \varphi(0) = 0, \varphi(T) = 0\}.$$

Definition 2.10. We say that a function $\theta_\varepsilon^\gamma \in \mathcal{V}$ is a corrector of order 0 iff

$$\left| \begin{array}{l} \varepsilon(\theta_\varepsilon^{\gamma'}, \varphi')_{L^2(Q)} + (\theta_\varepsilon^{\gamma'}, \varphi)_{L^2(Q)} + (\nabla \theta_\varepsilon^\gamma, \nabla \varphi)_{L^2(Q)} = \sqrt{\varepsilon} (f_\varepsilon, \varphi)_{L^2(Q)} \\ \theta_\varepsilon^\gamma(T) + y^\gamma(T) = 0, \end{array} \right. \quad \forall \mathcal{V}_0, \quad (33)$$

where we suppose that

$$\|f_\varepsilon\|_{L^2(]0,T[;H^{-1}(\Omega))} \leq C. \quad (34)$$

Getting a zero-order corrector We recall how to calculate a corrector of order 0 : We define $\varphi_\varepsilon^\gamma$ by writing

$$\begin{cases} -\varepsilon \varphi_\varepsilon^{\gamma''} + \varphi_\varepsilon^{\gamma'} & = 0, \\ \varphi_\varepsilon^\gamma(T) & = -y^\gamma(T), \end{cases}$$

$\varphi_\varepsilon^\gamma$ decreasing rapidly when $t \rightarrow -\infty$,

then

$$\theta_\varepsilon^\gamma(t) = -y^\gamma(T) e^{-\frac{T-t}{\varepsilon}}. \quad (35)$$

If we suppose that $y^\gamma(T) \in H_0^1(\Omega)$, the function

$$\theta_\varepsilon^\gamma = m \varphi_\varepsilon^\gamma \quad \left| \begin{array}{l} m = 1 \quad \text{in the neighbourhood of } t = T, \\ m = 0 \quad \text{in the neighbourhood of } t = 0 \end{array} \right.$$

is a corrector of order 0.

We then satisfy the variational equation, the main term being

$$m \Delta y^\gamma(T) e^{-\frac{T-t}{\varepsilon}} = \sqrt{\varepsilon} h_\varepsilon^\gamma.$$

Hence, under the above hypothesis we have :

$$\int_0^T \|h_\varepsilon^\gamma\|_{H^{-1}(\Omega)}^2 dt \leq C \varepsilon^{-1} \int_0^T e^{-\frac{2(T-t)}{\varepsilon}} dt = o(1).$$

We then have the theorem :

Theorem 2.11. Let be $\theta_\varepsilon^\gamma$ a corrector of order 0 defined by (33) and (34). We then have

$$\|y_\varepsilon^\gamma - (y^\gamma + \theta_\varepsilon^\gamma)\|_{L^2(]0,T[;H_0^1)} \leq C \sqrt{\varepsilon}. \quad (36)$$

Moreover,

$$\frac{d}{dt} [y_\varepsilon^\gamma - (y^\gamma + \theta_\varepsilon^\gamma)] \rightharpoonup 0 \quad \text{weakly in } L^2(Q), \quad (37)$$

when ε tends to 0.

Proof. If we put $w_\varepsilon^\gamma = y_\varepsilon^\gamma - (y^\gamma + \theta_\varepsilon^\gamma)$, then

$$\begin{aligned} & \varepsilon(w_\varepsilon^{\gamma'}, \varphi')_{L^2(Q)} + (w_\varepsilon^{\gamma'}, \varphi)_{L^2(Q)} + (\nabla w_\varepsilon^\gamma, \nabla \varphi)_{L^2(Q)} \\ & = -\varepsilon(y^{\gamma'}, \varphi')_{L^2(Q)} - \sqrt{\varepsilon} (f_\varepsilon, \varphi)_{L^2(Q)} \quad \forall \mathcal{V}_0. \end{aligned} \quad (38)$$

Particularly, if $\varphi = w_\varepsilon^\gamma$, then

$$\varepsilon \|w_\varepsilon^{\gamma'}\|_{L^2(Q)}^2 + \|w_\varepsilon^\gamma\|_{L^2(Q)}^2 \leq C \sqrt{\varepsilon} \left[\sqrt{\varepsilon} \|w_\varepsilon^{\gamma'}\|_{L^2(Q)} + \|w_\varepsilon^\gamma\|_{L^2(Q)} \right].$$

Thus the inequalities (36) and (37) hold.

2.3. Passage to the limit

We use the regularity properties of the heat equation, the *-well posed-* one, as follows :

First, we notice that $\|y_\gamma\|_{L^2(Q)} \leq C$ by the above proposition. It remains to see that ξ_γ (resp. p_γ) formally satisfies to the well-posed system :

$$(*) \left\{ \begin{array}{l} -\xi^{\gamma'} - \Delta \xi^\gamma = y^\gamma \in L^2(Q), \\ \xi^\gamma = 0, \\ \xi^\gamma(T) = 0, \end{array} \right. \quad (\text{resp. } (**)) \left\{ \begin{array}{l} -p^{\gamma'} - \Delta p^\gamma = y^\gamma - z_d, \\ p^\gamma = 0 \quad \text{on } \Sigma, \\ p^\gamma(T) = 0, \end{array} \right.$$

with the mean of a zero corrector. But (*) implies that

$$\|\xi^\gamma\|_{L^2(0,T; H_0^1(\Omega))} + \|\xi^{\gamma'}\|_{L^2(0,T; H^{-1}(\Omega))} \leq C,$$

(resp. (**)) gives $\|p^\gamma\|_{L^2(0,T; H_0^1(\Omega))} + \|p^{\gamma'}\|_{L^2(0,T; H^{-1}(\Omega))} \leq C$.

Then $\xi^\gamma \rightharpoonup \xi$ (resp. $p^\gamma \rightharpoonup p$) weakly in $L^2(0,T; H_0^1(\Omega))$, and by compactness $\xi^\gamma \rightarrow \xi$ (resp. $p^\gamma \rightarrow p$) strongly in $L^2(0,T; L^2(\Omega))$.

Also,

$$\left\{ \begin{array}{l} \rho^{\gamma'} - \Delta \rho^\gamma = 0, \\ \rho^\gamma = 0, \\ \rho^\gamma(0) = \lambda^\gamma(0), \end{array} \right.$$

implies that $\rho^\gamma \rightarrow \rho$ strong in $L^2(0,T; L^2(\Omega))$ by the same arguments, because

$$\|\lambda^\gamma(0)\|_{L^2(\Omega)} \leq \lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{\sqrt{\gamma}} \|\xi_\varepsilon'(u_\varepsilon^\gamma)(0)\|_{L^2(\Omega)} \leq C, \quad \text{then } \lambda^\gamma(0) \rightarrow \lambda(0) \in L^2(\Omega).$$

We then can announce the theorem :

Theorem 2.12. *The no-regret control u for the backward heat ill-posed problem (6), is characterized by the unique quadruplet $\{u, \xi, \rho, p\}$ solution to the optimality system :*

$$\left\{ \begin{array}{l} y' - \Delta y = u, \quad -\xi' - \Delta \xi = y \quad \rho' - \Delta \rho = 0, \\ \text{and} \quad -p' - \Delta p = y - z_d + \rho \quad \text{in } Q, \\ \\ y(0) = y(T) = 0, \quad \xi(T) = 0, \quad \text{and} \\ \rho(0) = \lambda(0), \quad p(T) = 0 \quad \text{in } \Omega, \\ \\ y = 0, \quad \xi = 0, \quad \rho = 0, \quad p = 0 \quad \text{on } \Sigma, \end{array} \right.$$

and the variational inequality :

$$\langle p + Nu, v - u \rangle \geq 0 \quad \forall v \in \mathcal{U}_{ad},$$

with $u \in \mathcal{U}_{ad}$, $y \in L^2(0,T; L^2(\Omega))$ and

$$p, \rho, \xi \in L^2(0,T; H^2(\Omega) \cap H_0^1(\Omega)), \quad \lambda \in L^2(\Omega).$$

Remark 2.13. As we have seen in this work, the hypothesis (12) is replaced by the no-regret notion. This method gives another point of view of solving the control problem of singular distributed systems.

3. THE CONTROLLABILITY APPROACH

Instead of the regularization for the backward ill-posed approach, we here use the null-controllability approach in Dorville *et al.* [3] for the regular heat equation (of incomplete data). With the low-regret method, we obtain at the end an optimality system for the backward ill-problem.

Consider the following heat system :

$$\begin{cases} \frac{\partial y}{\partial t} - \Delta y &= v + \theta \cdot \chi_\omega & \text{in } Q, \\ y &= 0 & \text{on } \Sigma, \\ y(0) &= g & \text{in } \Omega, \end{cases} \quad (39)$$

where $v \in L^2(Q)$, $\omega \subset\subset \Omega$ an open set, $g \in L^2(\Omega)$, and χ_ω such that

$$\chi_\omega u = \begin{cases} u & \text{on } (0, T) \times \omega, \\ 0 & \text{else} \end{cases}$$

For every fixed (v, g, θ) the system (39) admits a unique solution $y = y(v, g, \theta)$. The null controllability problem for (39) consists in finding $\theta \in L^2(0, T; L^2(\omega))$ such that if y is solution of (39) then

$$y(T) = 0. \quad (40)$$

This problem is classical and well-known (see for example Puel [16] and the references therein). Moreover, the solution is not unique.

Suppose selected a unique solution -by a given criterion- for (39)(40). More precisely, we will see that there is a criterion related to a weight function ρ such that for given (v, g) , there is a unique pair (θ, y_θ) such that

$$y_\theta = y(v, g; \theta) \quad \text{and} \quad y_\theta(T) = 0.$$

Denote by $\theta = \theta(v, g)$ and $y_\theta = y_\theta(v, g)$. We can notice that $y_\theta(0, 0) = 0$ iff $\theta(0, 0) = 0$. With this, for $(v, g) \in L^2(Q) \times L^2(Q)$, we consider the cost function

$$J(v, g) = \|y_\theta(v, g) - z_d\|_{L^2(Q)}^2 + N\|v\|_{L^2(Q)}^2 \quad (41)$$

where $z_d \in L^2(Q)$ and $N \in \mathbb{R}_+^*$ are given. Then the goal is to study the optimal control of this controlled system. The question of control for controllability problems of incomplete data is new, and was treated in detail by Dorville [1], and by Dorville *et al.* [3] very recently. We recall a variational method which ensures the existence and uniqueness selection of a solution for (39)(40).

The following lemma is slightly different (simpler) from lemma 1 in [3].

Lemma 3.1. Denote by $L = \frac{\partial}{\partial t} - \Delta$ and $L^* = -\frac{\partial}{\partial t} - \Delta$ the adjoint operator. Then there is a C^2 weighted positive function ρ on Q such that $\frac{1}{\rho}$ is bounded in Q , and there is a constant $C = C(\Omega, T, \omega, \rho) > 0$ such that

$$\int_Q \frac{1}{\rho^2} |q|^2 \, dxdt + \int_\Omega |q(0)|^2 \, dx \leq C \left[\int_Q |L^* q|^2 \, dxdt + \int_0^T \int_\omega \frac{1}{\rho^2} |q|^2 \, dxdt \right] \quad (42)$$

for every $q \in \mathcal{V} = \left\{ \varphi \in C^\infty([0, T] \times \overline{\Omega}) \mid \varphi|_\Sigma = 0 \right\}$.

This lemma holds from a Carleman inequality, and can be found in [4], or [15] and [16]. The RHS of (42) leads to use the following inner product :

$$a(r, s) = \int_Q L^* r L^* s \, dQ + \int_0^T \int_\omega \frac{1}{\rho^2} r s \, dQ. \quad (43)$$

Denote by $\|\cdot\|_a = \sqrt{a(\cdot, \cdot)}$ the associated norm on \mathcal{V} , and consider the completion given by the Hilbert space V . Finally, let be the linear form ℓ on V defined by

$$\ell(s) = \int_Q v s \, dQ + \int_\Omega g s(0) \, dx.$$

Proposition 3.2. *Under the hypothesis that $\|\rho v\|_{L^2(Q)} < +\infty$, there is a unique $\tilde{p} \in V$ solution to :*

$$a(\tilde{p}, s) = \ell(s) \quad \forall s \in V. \quad (44)$$

Moreover, if we put

$$y = L^* \tilde{p}, \quad \text{and} \quad \theta = -\frac{1}{\rho^2} \tilde{p} \cdot \chi_\omega, \quad (45)$$

then $\{y, \theta\}$ is the unique solution to (39)(40).

Proof. The proof follows from a simple application of the Lax-Milgram theorem, and by integration by parts.

Remark 3.3. Consider the set $L_\rho^2(Q) = \{w \in L^2(Q) \text{ such that } \rho w \in L^2(Q)\}$, and denote by $(\cdot, \cdot)_\rho = (\rho \cdot, \rho \cdot)_{L^2(Q)}$ and $\|\cdot\|_\rho = \sqrt{(\cdot, \cdot)_\rho}$. From proposition 3.2, for every pair (v, g) with $v \in L_\rho^2(Q)$ and $g \in L^2(\Omega)$, there is a unique pair $(\theta, y_\theta := y)$ solution of the problem (39)(40). We define the two linear functions

$$\begin{cases} L^2(Q) \times L^2(\Omega) & \longrightarrow & L^2(0, T; L^2(\omega)) \\ (v, g) & \longmapsto & \theta(v, g) \end{cases} \quad \text{and} \quad \begin{cases} L^2(Q) \times L^2(\Omega) & \longrightarrow & L^2(Q) \\ (v, g) & \longmapsto & y(v, g) \end{cases}$$

Then $v \mapsto \theta(v, 0) \in (L^2(0, T; L^2(\omega)), \rho dx dt)$ and $v \mapsto y(v, 0) \in L_\rho^2(Q)$ are continuous on $L_\rho^2(Q)$.

3.1. The low-regret control for the null-controllable problem

For every $v \in L_\rho^2(Q)$, we define the *weighted* cost function :

$$J_\rho(v, g) = \|y(v, g) - z_d\|_\rho^2 + N \|v\|_\rho^2 \quad (46)$$

where $z_d \in L_\rho^2(Q)$ and $N > 0$. Let be now \mathcal{U}_{ad}^ρ a closed convex subset of $L_\rho^2(Q)$.

By remark 3.3, we have $y(v, g) = y(v, 0) + y(0, g)$ and $\theta(v, g) = \theta(v, 0) + \theta(0, g)$. This, allows to write

$$J_\rho(v, g) - J_\rho(0, g) = J_\rho(v, 0) - J_\rho(0, 0) + 2(y(v, 0), y(0, g))_\rho.$$

Lemma 3.4. *Let be the operator $M : g \mapsto \tilde{p}(0, g)$ from $L^2(\Omega)$ to $\left(L^2(Q), \frac{1}{\rho} dx dt\right)$. Then M is linear continuous on $L^2(\Omega)$, and we have*

$$J_\rho(v, g) - J_\rho(0, g) = J_\rho(v, 0) - J_\rho(0, 0) + 2(S(v), g)_{L^2(\Omega)} \quad (47)$$

where S is a linear and continuous operator from $L^2_\rho(Q)$ to $L^2(\Omega)$, such that

$$S(v) = M^* (v + \theta(v, 0) \cdot \chi_\omega) + \tilde{p}(v, 0)(0), \quad (48)$$

where M^* is the adjoint of M .

Proof. The proof is simple and holds from (44) and remark 3.3. Indeed

$$\begin{aligned} (y(v, 0), y(0, g))_\rho &= \left(v - \frac{1}{\rho^2} \tilde{p}(v, 0) \cdot \chi_\omega, \tilde{p}(0, g) \right)_{L^2(Q)} + (\tilde{p}(v, 0), g)_{L^2(\Omega)} \\ &= \left(v - \frac{1}{\rho^2} \tilde{p}(v, 0) \cdot \chi_\omega, M g \right)_{L^2(Q)} + (\tilde{p}(v, 0), g)_{L^2(\Omega)}. \end{aligned}$$

We now give the optimality system of the low-regret control for the problem of missing data (39)-(40) and (46), defined by :

$$\inf_{v \in \mathcal{U}_{ad}^\rho} \left(\sup_{g \in L^2(\Omega)} \left(J_\rho(v, g) - J_\rho(0, g) - \gamma \|g\|_{L^2(\Omega)}^2 \right) \right) \quad (49)$$

for $\gamma > 0$. Using (47), the problem is equivalent to the following classical control problem

$$\inf_{v \in L^2_\rho(Q)} \mathcal{J}_\rho^\gamma(v) \quad \text{where} \quad \mathcal{J}_\rho^\gamma(v) = J_\rho(v, 0) - J_\rho(0, 0) + \frac{1}{\gamma} \|S(v)\|_{L^2(\Omega)}^2. \quad (50)$$

Lemma 3.5. *The problem (50) admits a unique solution $u_\gamma \in \mathcal{U}_{ad}^\rho$.*

Proof. We have the following :

$\mathcal{J}_\rho^\gamma(v) \geq -J_\rho(0, 0) \forall v \in \mathcal{U}_{ad}^\rho$. Hence $d_\gamma = \inf_{v \in \mathcal{U}_{ad}^\rho} \mathcal{J}_\rho^\gamma(v)$ exists. Let then $v_n = v_n(\gamma)$ be a minimizing subsequence such that $d_\gamma = \lim_{n \rightarrow \infty} \mathcal{J}_\rho^\gamma(v_n)$. We have

$$-J_\rho(0, 0) \leq \mathcal{J}_\rho^\gamma(v_n) = J_\rho(v_n, 0) - J_\rho(0, 0) + \frac{1}{\gamma} \|S(v_n)\|_{L^2(\Omega)}^2 \leq d_\gamma + 1.$$

We deduce the following estimates :

$$\|v_n\|_\rho \leq c_\gamma, \quad \frac{1}{\sqrt{\gamma}} \|S(v_n)\|_{L^2(\Omega)} \leq c_\gamma, \quad \|y(v_n, 0) - z_d\|_\rho \leq c_\gamma,$$

(c_γ independent of n). Then, there is $u_\gamma \in \mathcal{U}_{ad}^\rho$ such that $v_n \rightharpoonup u_\gamma$ weak in $L^2_\rho(Q)$. Also, $y(v_n, 0) \rightharpoonup y(u_\gamma, 0)$ (continuity with respect to the data). At last, thanks to the convexity of \mathcal{J}_ρ^γ , we deduce that u_γ is unique.

Theorem 3.6. *The low-regret control $u_\gamma \in \mathcal{U}_{ad}^\rho$ is characterized by the quadruplet $\{u_\gamma, y_\gamma, \tilde{\theta}_\gamma, p_\gamma\} \in \mathcal{U}_{ad}^\rho \times L^2(Q) \times V \times L^2(Q)$ unique solution of the system*

$$\begin{cases} L y_\gamma &= u_\gamma + \theta(u_\gamma, 0) \cdot \chi_\omega, & L^* p_\gamma &= y_\gamma - z_d + \tilde{\theta}_\gamma \cdot \chi_\omega & \text{in } Q, \\ y_\gamma &= 0, & p_\gamma &= 0 & \text{on } \Sigma, \\ y_\gamma(0) = p_\gamma(0) &= 0, & y_\gamma(T) = p_\gamma(T) &= 0 & \text{in } \Omega, \end{cases} \quad (51)$$

where $y_\gamma = y(u_\gamma, 0)$, $p_\gamma = p(u_\gamma, 0)$, $\tilde{\theta}_\gamma = \tilde{\theta}(u_\gamma, 0)$, and the variational inequality

$$\left(T^* [L p_\gamma - \tilde{\theta}_\gamma \cdot \chi_\omega] + N \rho^2 u_\gamma + (1/\gamma) S^* S(u_\gamma), v - u_\gamma \right)_{L^2(Q)} \geq 0 \quad \forall v \in \mathcal{U}_{ad}^\rho \quad (52)$$

where S^* is the adjoint of S , and T^* is the adjoint of the operator $T : v \mapsto L^* \tilde{p}(v, 0)$.

Proof. The necessary condition of Euler-Lagrange satisfied by u_γ gives

$$(y_\gamma - z_d, \rho^2 y(v - u_\gamma, 0))_{L^2(Q)} + N(\rho^2 u_\gamma, v - u_\gamma)_{L^2(Q)} + \frac{1}{\gamma} (S(u_\gamma), S(v - u_\gamma))_{L^2(\Omega)} \geq 0 \quad \forall v \in \mathcal{U}_{ad}^\rho,$$

$y_\gamma = y(u_\gamma, 0)$. We then introduce $\sigma_\gamma = \sigma(u_\gamma, 0) \in V$, unique solution of the problem

$$a(\sigma_\gamma, q) = \int_Q (y_\gamma - z_d) q \, dxdt, \quad \forall q \in V.$$

We then define the *adjoint* state $p_\gamma = p(u_\gamma, 0)$ by

$$p_\gamma = L^* \sigma_\gamma \quad \text{and we put} \quad \tilde{\theta}_\gamma = -\frac{1}{\rho^2} \sigma_\gamma.$$

Then the pair $(\tilde{\theta}_\gamma, p_\gamma)$ is solution of the null-controllability problem :

$$\left\{ \begin{array}{ll} L^* p_\gamma & = y_\gamma - z_d + \tilde{\theta}_\gamma \cdot \chi_\omega \quad \text{in } Q, \\ p_\gamma & = 0 \quad \text{on } \Sigma, \\ p_\gamma(0) = p_\gamma(T) & = 0 \quad \text{in } \Omega \end{array} \right.$$

Then the Euler condition writes :

$$(Lp_\gamma - \tilde{\theta}_\gamma \cdot \chi_\omega, \rho^2 y(v - u_\gamma, 0))_{L^2(Q)} + (N\rho^2 u_\gamma + (1/\gamma)S^*S(u_\gamma), v - u_\gamma)_{L^2(Q)} \geq 0.$$

From another side we have

$$(Lp_\gamma - \tilde{\theta}_\gamma \cdot \chi_\omega, \rho^2 y(v - u_\gamma, 0))_{L^2(Q)} = (T^*(Lp_\gamma - \tilde{\theta}_\gamma \cdot \chi_\omega), v - u_\gamma)_{L^2(Q)} \quad \forall v \in \mathcal{U}_{ad}^\rho,$$

where T^* is the adjoint of the linear continuous operator $T : v \mapsto L^* \tilde{p}(v, 0)$ de $L_\rho^2(Q)$ in $(L_\rho^2(Q), \rho^2 \, dxdt)$.

3.2. Passage to the limit

In this section, we verify that we also have an optimality for the no-regret control. It remains to see that y_γ (resp. p_γ) particularly satisfies to the well-posed system :

$$(3^*) \left\{ \begin{array}{l} Ly_\gamma = u_\gamma + \theta(u_\gamma, 0)\chi_\omega \in L^2(Q), \\ y_\gamma = 0, \\ y_\gamma(0) = 0, \end{array} \right. \quad (\text{resp. } (4^*) \left\{ \begin{array}{l} L^* p_\gamma = y_\gamma - z_d + \tilde{\theta}_\gamma \chi_\omega, \\ p_\gamma = 0 \quad \text{on } \Sigma, \\ p_\gamma(T) = 0, \end{array} \right.)$$

and of course $y_\gamma(T) = 0$ (resp. $p_\gamma(0) = 0$).

The system (3*) implies

$$\|y_\gamma\|_{L^2(0,T; H_0^1(\Omega))} + \|y_\gamma'\|_{L^2(0,T; H^{-1}(\Omega))} \leq C,$$

(resp. (4*) gives $\|p_\gamma\|_{L^2(0,T; H_0^1(\Omega))} + \|p_\gamma'\|_{L^2(0,T; H^{-1}(\Omega))} \leq C$).

Then $y_\gamma \rightharpoonup y$ (resp. $p_\gamma \rightharpoonup p$) weakly in $L^2(0, T; H_0^1(\Omega))$, and by compactness $y_\gamma \rightarrow y$ (resp. $p_\gamma \rightarrow p$) strongly in $L^2(0, T; L^2(\Omega))$.

Also,

$$\frac{1}{\gamma} S^* S(u_\gamma) \rightarrow S^* S(u)$$

strongly in $L^2(0, T; L^2(\Omega))$ by the same arguments, since

$$\|S^*S(u)\|_{L^2(\Omega)} \leq \lim_{\gamma \rightarrow 0} \frac{1}{\sqrt{\gamma}} \|S^*S(u_\gamma)\|_{L^2(\Omega)} \leq C.$$

We then obtain the following singular optimality system, stated in the following theorem :

Theorem 3.7. *The no-regret control $u \in \mathcal{U}_{ad}^p$ is characterized by the quadruplet $\{u, y, \tilde{\theta}, p\} \in \mathcal{U}_{ad}^p \times L^2(Q) \times V \times L^2(Q)$, unique solution of the system*

$$\begin{cases} Ly &= u + \theta(u, 0) \cdot \chi_\omega, & L^*p &= y - z_d + \tilde{\theta} \cdot \chi_\omega & \text{in } Q, \\ y &= 0, & p &= 0 & \text{on } \Sigma, \\ y(0) = p(0) &= 0, & y(T) = p(T) &= 0 & \text{in } \Omega, \end{cases} \quad (53)$$

where $y = y(u, 0)$, $p = p(u, 0)$, $\tilde{\theta} = \tilde{\theta}(u, 0)$,
with the variational inequality

$$\left(T^*[Lp - \tilde{\theta} \cdot \chi_\omega] + N\rho^2 u + S^*S(u), v - u \right)_{L^2(Q)} \geq 0 \quad \forall v \in \mathcal{U}_{ad}^p, \quad (54)$$

where S^* is the adjoint of S , and T^* the adjoint of the operator $T : v \mapsto L^*\tilde{p}(v, 0)$.

Remark 3.8. As a general remark on this section, we see that the optimality system has the same structure as the one of standard optimal control problems. More precisely, the adjoint state satisfies to a null-controllability system, as for the state y .

REFERENCES

- [1] Dorville R. *Sur le contrôle de quelques problèmes singuliers associés à l'équation de la chaleur (On the control of some singular problems associated to the heat equation)*. Thèse de Doctorat de l'Université des Antilles et de la Guyane, Guadeloupe (2004).
- [2] Dorville R., Nakoulima O., Omrane A. *Low-regret control for singular distributed systems: The backwards heat ill-posed problem*. Applied Math. Letters, 17/5, pp 549–552 (2004).
- [3] Dorville R., Nakoulima O., Omrane A. *Contrôle optimal pour les problèmes de contrôlabilité de systèmes distribués à données manquantes*. C. R. Acad. Sci. Paris Ser. I Math., Vol. 338, pp 921–924 (2004).
- [4] Emanuilov O. Yu. *Controllability of Parabolic Equations*. Sbornik Mathematics, Vol. 186, No 6, pp 879–900 (1995).
- [5] Lions J. L. *Contrôle optimal des systèmes gouvernés par des équations aux dérivées partielles*. Dunod, Paris (1969).
- [6] Lions J. L. *Contrôle optimal pour les systèmes distribués singuliers*. Gauthiers-Villard, Paris (1983) .
- [7] Lions J. L. *Perturbations singulières dans les problèmes aux limites et en contrôle optimal*. Lecture Notes in Mathematics, Springer (1973).
- [8] Lions J. L. *Contrôle à moindres regrets des systèmes distribués*. C. R. Acad. Sci. Paris Ser. I Math., Vol. 315, pp 1253–1257 (1992).
- [9] Lions J. L. *No-regret and low-regret control*, Environment, Economics and Their Mathematical Models, Masson, Paris, 1994.
- [10] Lions J. L. *Duality Arguments for Multi Agents Least-Regret Control*. Collège de France, Paris (1999).
- [11] Lions J.-L., Magenes E. *Problèmes aux limites non homogènes (I) et (II)*, Ann. Sc. Norm. Sup. Pisa, Vol. 14, pp 259–308, et Vol. 15, (1962), pp 39–101 (1961) .
- [12] Lions J.-L., Magenes E. *Remarques sur les problèmes aux limites linéaires elliptiques*, Rend. Acc. Naz. Lincei, Vol. 8/32, pp 873–883 (1962).
- [13] Nakoulima O., Omrane A., Velin J. *Perturbations à moindres regrets dans les systèmes distribués à données manquantes*. C. R. Acad. Sci. Paris Ser. I Math., Vol. 330, pp 801–806 (2000).
- [14] Nakoulima O., Omrane A., Velin J. *On the pareto control and no-regret control for distributed systems with incomplete data*. SIAM J. Control Optim. vol. 42, N 4, pp. 1167–1184 (2003).
- [15] Puel J. P. *Contrôlabilité exacte et approchée*. Cours de DEA, Université de Paris 6 (2000).
- [16] Puel J. P. *Applications of global Carleman inequalities to controllability and inverse problems*. (to appear)
- [17] Savage L. J. *The Foundations of Statistics*. Dover, 2nd Edition (1972).