



HAL
open science

An holistic approach to cropping practices, a prerequisite for an agroecological diagnosis of plantain cropping systems in Guadeloupe.

Claire Forite, Alexandre Ogisma, Brunise Deloné-Louis-Jeune, Gladys Loranger-Merciris, Harry Ozier-Lafontaine, Jean-Louis Diman

► To cite this version:

Claire Forite, Alexandre Ogisma, Brunise Deloné-Louis-Jeune, Gladys Loranger-Merciris, Harry Ozier-Lafontaine, et al.. An holistic approach to cropping practices, a prerequisite for an agroecological diagnosis of plantain cropping systems in Guadeloupe.. 47th Annual Scientific Meeting of the Caribbean Food Crops Society, Jul 2011, Lloyd Erskine Sandiford Centre, Bridgetown, Barbados. hal-04089246

HAL Id: hal-04089246

<https://hal.univ-antilles.fr/hal-04089246v1>


Submitted on 13 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An holistic approach to cropping practices, a prerequisite for an agroecological diagnosis of plantain cropping systems in Guadeloupe.

Claire Forite^{1,2}, Alexandre Ogisma^{1,3,4}, Brunise Delone-Louis-Jeune^{3,5}, Gladys Loranger-Merciris^{1,3}, Harry Ozier-Lafontaine¹, Jean-Louis Diman¹



- Context: No studies and no technical support for farmers about plantain production in Guadeloupe
Plantain cropping systems are mainly improved in the Cavendish production area
Cropping system duration is reduced to 1 or 2 years long because of telluric pathogens' pressure
- Aim: Characterize plantain cropping system diversity in order to run an agroecological diagnosis
Identify the factors influencing the cropping system sustainability
- Hypothesis: Plantain cropping practices are influenced by Cavendish reference cropping system


METHOD

- Survey improvement
- Plantain cropping systems typology
- Selection of 2 contrasted agroecological situations
- Plots' agroecological diagnosis



PLANTAIN CROPPING SYSTEM TYPOLOGY

Different levels of Cavendish export cropping management deviation depending on incomes and plant management




Selection of plots from type 1 vs type 4

AGROECOLOGICAL DIAGNOSIS FOR CONTRASTED SITUATIONS

-Yield measures


Number of bananas on the bunch
fonction of pseudostem circumference




- Type 1: higher yields than type 4
- Type 1: better homogeneity of bunches in "blanc" variety
- Type 4: "Blanc" variety seems to be more productive

- Higher nematodes' infestation in type 1
- Different nematodes species combination in type 1 & 4
- Less roots' necroses in type 1


- Nematodes identification and counting



-Weevils infestation image analyzing



- Leaf diagnostic



CONCLUSION

- Different cropping systems according to the production area (type 4: out of the main Cavendish production area)
- Higher yields in more intensive systems

REMAINING QUESTIONS: What are the investments and benefits for each cropping system? What is cropping systems sustainability?

Caribbean Agroecology Networking Symposium

"Productive Agrosystems and Resources Conservation in an Island Environment: an Agroecological Challenge for Caribbean Sustainable Development"
47th Annual Scientific Meeting of the Caribbean Food Crops Society, Lloyd Erskine Sandiford Centre, Bridgetown, Barbados, July 3-9, 2011

