



**HAL**  
open science

# Skin Application of 4% Menthol Enhances Running Performance in Hot and Humid Climate

Eric Hermand, Aurelie Collado, Olivier Hue

► **To cite this version:**

Eric Hermand, Aurelie Collado, Olivier Hue. Skin Application of 4% Menthol Enhances Running Performance in Hot and Humid Climate. *International Journal of Sports Medicine*, 2020, 41 (03), pp.161-167. 10.1055/a-1062-6520 . hal-04585897

**HAL Id: hal-04585897**

**<https://hal.univ-antilles.fr/hal-04585897v1>**

Submitted on 27 May 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Skin Application of 4% Menthol Enhances Running Performance in Hot and Humid Climate

## Authors

Eric Hermand<sup>1,2,3</sup> , Aurelie Collado<sup>2</sup>, Olivier Hue<sup>2</sup>

## Affiliations

- 1 EA 6310 'HAVAÉ': Handicap, Activité, Vieillesse, Autonomie, Exercice, Limoges University, Limoges, France
- 2 EA 3596 'ACTES': Adaptation, Climat Tropical, Exercice et Santé, Université des Antilles et de la Guyane, Pointe-a-Pitre, Guadeloupe
- 3 UMR INSERM U1272: Hypoxie & Poumon, Paris 13 University - Bobigny Campus, Bobigny, France

## Key words

tropical climate, aerobic performance, running, menthol, external cooling

accepted 03.11.2019

## Bibliography

DOI <https://doi.org/10.1055/a-1062-6520>

Int J Sports Med 2019; 40: 1–7

© Georg Thieme Verlag KG Stuttgart · New York

ISSN 0172-4622

## Correspondence

Dr. Eric Hermand

Université des Antilles et de la Guyane, Laboratoire ACTES

Campus Fouillole

STAPS

97157 Pointe-a-Pitre

Guadeloupe

Tel.: 0590/48/31/71, Fax : 0590/48/31/79

[eric.hermand17@gmail.com](mailto:eric.hermand17@gmail.com)

## ABSTRACT

Aerobic performance is negatively impacted by tropical climate due to impairment of thermoregulatory mechanisms. We tested the hypothesis that a torso application of a 4% menthol solution would have the same effect on a best performance 10-km run as an external use of cold water. Thirteen trained male athletes completed four outdoor 10-km runs ( $T = 29.0 \pm 1.3^\circ\text{C}$ , relative humidity  $59.0 \pm 13.6\%$ ) wearing a tee-shirt soaked every 2-km either in a cold ( $\sim 6^\circ\text{C}$ ) or warm/ambient ( $\sim 28^\circ\text{C}$ ) solution, consisting in water or in a 4% menthol solution, (CTL, MENT-Amb, CLD and MENT-CLD). Run performances were improved from 4.8 to 6.1% in CLD ( $51.4 \pm 5.5$  min), MENT-Amb ( $52.2 \pm 5.9$  min) and MENT-CLD ( $51.4 \pm 5.1$  min) conditions (vs. CTL,  $55.4 \pm 8.4$  min,  $P < 0.05$ ), without differences between these three conditions, whereas heart rate ( $177 \pm 13$  bpm), body temperature ( $38.7 \pm 0.6^\circ\text{C}$ ) and drink ingestion ( $356 \pm 170$  g) were not modified. Thermal sensation after running was lower in MENT-CLD (vs. CTL,  $P < 0.01$ ) and thermal acceptability was higher in CLD and MENT-Amb (vs. CTL,  $P < 0.05$ ), but thermal comfort, feeling scale and rate of perceived exertion remained unchanged. The use of menthol on skin enhances aerobic performance in a tropical climate, and no differences in performance were observed between menthol and traditional percooling strategies. However, combining both menthol and traditional percooling brought no further improvements.

## Introduction

A hot and humid climate decreases physical performance, especially in aerobic sports [1]. Although the mechanisms leading to these alterations are not fully understood, it is well established that thermoregulatory system dysfunction is involved with the loss of ability to dissipate excess metabolic heat [2]. These mechanisms, already impaired in a hot environment, are further altered in humid climate, which drastically limits evapotranspiration due to the water-saturated atmosphere [3]. Hence, running may be the sport most impacted by a tropical climate because heat dissipation by circulating air convection around the body, such as in cycling, is reduced [4]. To limit its effect in endurance sports and to enhance

thermoregulatory processes, several countermeasures have been studied with various outcomes on performance: acclimation, hydration, pacing, and cooling strategies [1].

Among the latter, external cooling solutions exist in the form of pads containing low-temperature inserts that may be applied on different body parts such as the arms, legs, torso, and neck [5]. Used before (pre-) or during (per-) exercise, they allow better performance in hot and humid environment by limiting the rise of body core temperature [6]. For example, performances from sprints to short-distance running (up to 5 km) are enhanced by wearing 'ice vests' during warm-up [7], and the use of a cooling neck collar augments the limit time to exhaustion at aerobic submaximal intensi-

■ Proof copy for correction only. All forms of publication, duplication or distribution prohibited under copyright law. ■

ty [8]. Performances are further improved when methods are combined (pre- and per-) [9] and/or used with a more “aggressive” cooling temperature of around 0 °C [6]. Indeed, the disequilibrium between heat production and heat loss during prolonged exercise, leading to hyperthermia, has a negative impact on physiological functions and exercise performance: heat-dissipating mechanisms (skin vasodilatation, sweating response) are not enough to maintain an optimal body core temperature. Strategies relying on pre- (before exercise) and percooling (during) then augment heat storage and diminish the thermoregulatory strain.

Beyond traditional approaches of body cooling, which are efficient in laboratory-controlled environment but difficult to use in ecological conditions, alternative strategies have recently been developed to maximize performance in tropical conditions. One of them is menthol, a compound largely used in the food and tobacco industries, and popular in Asia for its therapeutic properties [10]. It mostly acts on TRPM8 and TRPA1 thermoreceptors [11] on the skin and internal mucous membranes. When activated by menthol on the skin, they induce a local cold sensation that is prone to modify thermal sensation, especially in hot environment, [12], which leads to reduced fatigue perception during maximal and submaximal exercise, and potentially to better performance [13, 14]. These preliminary observations should have led to further comparisons between the respective uses of menthol and proven external cooling. However, to our knowledge, there is no study directly comparing the use of menthol at ambient temperature on skin and traditional cooling strategies to test if menthol could allow the same level of performance in an ecologically hot and humid environment without the equipment burden. In other conditions, in a laboratory-controlled environment (34 °C, RH 30%), there was no difference of performance between the use of cold 4% menthol gel on the skin and cold pads, both applied around the heat-sensitive neck compared to control condition [15]. Barwood et al. showed no difference in running performance between sprayed 0.05% menthol and water at 20 °C [16].

Hence, in view of the lack of data comparing skin-applied menthol and cold-water cooling, because aerobic performance is potentially reduced by the neurosensitive perception of environmental heat and humidity, we hypothesize that the use of menthol on skin would blunt these perception inhibition mechanisms and lead to enhanced performances, similar to the external use of cold water for body cooling.

## Material and Methods

### Subjects

Thirteen moderately to well-trained and heat-acclimatized male athletes, with a performance level from 3–4 out of 5 [17], participated in this study (age: 21 ± 4 years, height: 176 ± 6 cm, body mass: 70 ± 9 kg, maximal aerobic speed 16.2 ± 1.3 km·h<sup>-1</sup>). All participants were subject to a prior medical examination to check for any cardiopulmonary disease or previous heat stroke, and were tested for their maximal aerobic speed [18]. The study was approved by the National Ethic Committee (CPP, registration number 2018-A00295–50) and subjects gave their informed consent. Procedures conformed with ethics in sport and exercise science [19].

### Experimental design

The protocol was conducted in ecological conditions: participants in this field study ran outside, on a flat paved road, directly exposed to the tropical heat and humidity of the French West Indies (wet-bulb globe temperature (WBGT): 29.0 ± 1.3 °C, relative humidity 59.0 ± 13.6%), such as any athletes would face in endurance races held in the same environment.

Subjects completed four 10-km run tests as fast as they could on a flat course, 4–7 days apart, in a randomized cross-over design including four experimental conditions in which the athlete’s shirt would be soaked in four different solutions: (1) control (CTL): water at ambient temperature (T = 28.7 ± 2.9 °C); (2) cold water (CLD) at low temperature (T = 6.0 ± 0.8 °C); (3) menthol at ambient temperature (MENT-Amb): 4% menthol solution at ambient temperature (T = 28.2 ± 2.3 °C); and (4) menthol at low temperature (MENT-CLD): 4% menthol solution at cold temperature (T = 6.1 ± 0.6 °C). The menthol solutions were prepared from an 86% menthol-concentrated menthol solution (Robertet, Grasse, France), which was diluted to obtain a 4% solution. Athletes were asked to restrain from training the day before the tests and avoid caffeine on test day. For every athlete, trials were undertaken at the same time of the day to limit both variations of wet-bulb globe temperature (WBGT) throughout the day and circadian variations of core temperature. Finally, athletes were asked to wear the same outfit at all sessions, except the white shirt provided by the experimenter.

### Experimental procedure

Heart rate (HR), stride rate, and run duration were recorded continuously (sampling frequency 1 s) using an M400 Polar watch paired with an H7 strap belt or an OH1 sensor [20, 21] (Polar Electro Oy, Kempele, Finland). WBGT was continuously measured during the trials with a HD32.2 device (Delta Ohm, Padova, Italia), placed immediately next to the run course. Core temperature (T<sub>co</sub>) was telemetrically measured via ingestible temperature measurement pills (BodyCap, Caen, France), with an embedded memory (sampling frequency 30 s). Athletes were instructed to ingest these pills 6 to 8 h before trials to ensure the pill was out of the stomach, thereby avoiding changes in T<sub>co</sub> due to fluid consumption. Each trial included a preliminary 15- to 20-min standardized warm-up. After 5 min of gear application and weighing, subjects started the 10-km run on the 1-km out-and-back course. At the start, and every 2 km, athletes stopped for 30–40 s, during which: 1) athletes took their shirt off; 2) the experimenter soaked it in a solution corresponding to the tested condition (CTL, CLD, MENT-Amb or MENT-CLD), adding or renewing between 250 and 300 g of solution in T-shirt fabric; 3) athletes put back their shirt on; 4) athletes could hydrate ad libitum with water at ambient temperature (T = 26.9 ± 1.6 °C). The 2-km stops were necessary due to the pronounced sweating rate in the hot and humid condition, which would drip the solution off the shirt and blunt its potential effects. Finally, psychological parameters were assessed via oral or written questionnaires immediately before the start: feeling scale (FS, from –5 ‘Very bad to +5 ‘Very good’) [22]; thermal comfort (TC, from –3 ‘Very uncomfortable’ to +3 ‘Very comfortable’) [23]; thermal sensation (TS, from –3 ‘Very cold’ to +3 ‘Very hot’) [24]; thermal acceptability (TA, from –1 ‘Clearly unacceptable’ to +1 ‘Clearly acceptable’) [25]; and

post-race FS, TC, TS, TA, and rate of perceived exertion (RPE, from 6 'Very, very light' to 20 'Very, very hard') [26–28].

## Data collection and statistical analyses

Following the guidelines of the National Ethics Committee, the different run conditions were strictly randomized to avoid effects of trial order. Run duration, HR, and stride rate data were measured and averaged on the successive 2-km intervals and on the whole run, and were retrieved from the Polar platform website.  $T_{co}$  values were telemetrically updated and downloaded at the end of each session from monitors. Using calibrated balances (Terraillon, Croissy-sur-Seine, France), athletes were weighed-in, dry, shoes and shirt off, immediately before and after each run and water consumption (g) was measured after each stop. Water loss during the run was then calculated by adding the weight variation to the quantity of consumed water, and the percentage of weight loss was extracted using initial body weight. Psychological parameters were compiled before and after each run. Normality of data on each condition was verified by a Kolmogorov-Smirnov test. Thus, repeated measures ANOVAs were conducted with Condition (i. e., CTL, CLD, MENT-Amb, MENT-CLD) as the between-factor and Time (i. e., 2-km intervals) as the within-factor. Posthoc Student's t-tests were performed, when applicable, to compare mean values between successive intervals ([0–2 km] vs. [2–4 km], [2–4 km] vs. [4–6 km], [4–6 km] vs. [6–8 km] and [6–8 km] vs. [8–10 km]) to assess the kinetics of the run pace throughout the 10-km course and between conditions. Finally, the effect size was assessed by computing Hedge's *g* when applicable. Statistical significance was set at  $P < 0.05$ .

## Results

Results are presented in ► **Table 1** and in ► **Fig. 1** and ► **2**.

► **Table 1** Mean values ( $\pm$ SD) of environmental condition (WBGT and RH); RPE from 6 'Very, very light' to 20 'Very, very hard'; FS from –5 'Very bad to +5 'Very good; TS from –3 'Very cold' to +3 'Very hot'; TC from –3 'Very uncomfortable' to +3 'Very comfortable'; TA from –1 'Clearly unacceptable' to +1 'Clearly acceptable'; ingested drink and percentage of weight loss.

		CTL	CLD	MENT-AMB	MENT-CLD
<b>Environmental conditions</b>	WBGT (°C)	29.1 $\pm$ 1.5	28.9 $\pm$ 1.2	28.8 $\pm$ 1.5	29.1 $\pm$ 1.2
	RH (%)	57.7 $\pm$ 12.9	60.0 $\pm$ 13.4	63.6 $\pm$ 14.5	54.6 $\pm$ 13.3
<b>RPE (n.u.)</b>		15.8 $\pm$ 2.2	14.5 $\pm$ 3.1	14.6 $\pm$ 1.8	15.6 $\pm$ 1.7
<b>FS (n.u.)</b>	Pre	2.0 $\pm$ 2.0	2.3 $\pm$ 2.1	2.5 $\pm$ 2.1	1.9 $\pm$ 2.3
	Post	–0.8 $\pm$ 2.3	0.1 $\pm$ 2.3	0.7 $\pm$ 2.0 *	–0.9 $\pm$ 1.7 †
<b>TS (n.u.)</b>	Pre	1.2 $\pm$ 0.9	0.7 $\pm$ 0.9	1.0 $\pm$ 0.8	1.1 $\pm$ 1.0
	Post	1.2 $\pm$ 1.4	1.1 $\pm$ 0.9	0.0 $\pm$ 1.7	–0.4 $\pm$ 1.2 * * * + +
<b>TC (n.u.)</b>	Pre	0.5 $\pm$ 0.7	0.6 $\pm$ 0.6	0.8 $\pm$ 0.7	0.6 $\pm$ 0.8
	Post	–0.5 $\pm$ 1.2	–0.1 $\pm$ 1.2	0.3 $\pm$ 0.9	0.2 $\pm$ 0.6
<b>TA (n.u.)</b>	Pre	0.3 $\pm$ 0.4	0.3 $\pm$ 0.3	0.5 $\pm$ 0.3	0.3 $\pm$ 0.4
	Post	–0.1 $\pm$ 0.5	0.2 $\pm$ 0.4 *	0.4 $\pm$ 0.5 *	0.2 $\pm$ 0.5
<b>Drink ingested (g)</b>		363 $\pm$ 173	367 $\pm$ 162	318 $\pm$ 164	379 $\pm$ 198
<b>% weight loss</b>		–1.3 $\pm$ 0.8	–1.0 $\pm$ 0.6	–1.4 $\pm$ 1.0	–1.2 $\pm$ 0.6

SD, standard deviation; WBGT, wet-bulb globe temperature; RH, relative humidity; RPE, rate of perceived exertion; FS, feeling scale; TS, thermal sensation; TC, thermal comfort; TA, thermal acceptability; CTL, control; CLD, cold; MENT-AMB, menthol at ambient temperature; MENT-CLD, menthol at low temperature; n.u., no unit or normalized unit. \*  $P < 0.05$ , \* \*  $P < 0.01$ : vs. CTL. + +  $P < 0.01$ : vs. CLD. †  $P < 0.05$ : vs. MENT-AMB

## Environmental conditions

WBGT and relative humidity are presented in ► **Table 1**. There was no difference between conditions.

## 10-km performance (► **Fig. 1a**)

Athletes were faster in CLD (–6.1 %), MENT-Amb (–4.8 %), and MENT-CLD (–6.1 %) when compared with control ( $P < 0.05$ ,  $g = 0.55$ , 0.44, 0.57 respectively). However, no differences were observed between MENT-Amb, MENT-CLD, and CLD (► **Fig. 1a**).

## 2-km intervals

Interval splits (i. e., running performance) increased for all conditions ( $P < 0.01$ , ► **Fig. 1a**), meaning that athletes were getting slower throughout the run.

There was a condition effect on 2-km splits ( $P < 0.01$ , ► **Fig. 1a**). No differences were observed between conditions in the first two intervals. From the [4–6 km] interval to the last, split durations were longer in CTL (vs. CLD, Ment-Amb, and Ment-CLD,  $P < 0.05$ , ► **Fig. 1a** and **b**).


Interval splits continuously increased from the start in the CTL condition ([0–2 km] vs. [2–4 km],  $P < 0.05$ ; [2–4 km] vs. [4–6 km],  $P < 0.001$ ; [4–6 km] vs. [6–8 km],  $P < 0.01$ ) before stabilizing during the two last intervals. In CLD, Ment-Amb, and Ment-CLD, after some pace variations (or not), speed was not modified on the last three intervals ([4–6 km], [6–8 km], and [8–10 km]).

## Heart rate


No differences were observed between conditions (► **Fig. 1c** and **d**).

## Stride rate

Stride rate was not modified by condition (► **Fig. 2a** and **b**).



► **Fig. 1** a and b overall run duration in CTL, CLD, MENT-Amb, and MENT-CLD conditions a, upper panel: individual values and mean ± SD) and by 2 km-interval per condition for each condition b, lower panel: mean ± SEM). c and d average heart rate in CTL, CLD, MENT-Amb, and MENT-CLD conditions c, upper panel: mean ± SD) and by 2 km-interval per condition for each condition d, lower panel: mean ± SEM). \*: P < 0.05, CLD/MENT-Amb/MENT-CLD vs. CTL. Time effects were not reported for readability.



► **Fig. 2** a and b average stride rate in CTL, CLD, MENT-Amb, and MENT-CLD conditions a, upper panel: mean ± SD) and by 2 km-interval per condition for each condition b, lower panel: mean ± SEM). c and d average core temperature in CTL, CLD, MENT-Amb, and MENT-CLD conditions c, upper panel: mean ± SD) and by 2 km-interval per condition for each condition d, lower panel: mean ± SEM). Time effects were not reported for readability.

## Core temperature

$T_{co}$  increased throughout the run ( $P < 0.001$ ) for each condition (► **Fig. 2d**,  $P < 0.001$ ). However, there was no difference between conditions (► **Fig. 2c**).

## Perceptual measures

At the end of the run, FS, TS, TC, and TA were all lower ( $P < 0.001$  and  $g = 1.12$ ,  $P < 0.05$  and  $g = 0.40$ ,  $P < 0.001$  and  $g = 0.71$ , and  $P < 0.05$  and  $g = 0.44$ , respectively) compared to the trial start. TS at the end of the run was lower in MENT-CLD vs. CTL, and vs. CLD ( $P < 0.01$ ,  $g = 1.23$  and  $1.22$  respectively). Similarly, TA was increased in CLD, MENT-Amb (vs. CTL,  $P < 0.05$ ,  $g = 0.74$  and  $1.04$  respectively) (► **Table 1**). Condition had an effect on TA ( $P < 0.05$ , ► **Table 1**). RPE, FS, and TC were not modified by conditions (► **Table 1**).

## Drink ingestion (► **Table 1**)

Fluid ingestion increased throughout the run ( $P < 0.01$ ), although there was no difference in fluid ingestion between immediate successive intervals. However, some differences were observed between the [0–2 km] and [4–6 km] intervals (CLD,  $P < 0.01$ ; MENT-CLD,  $P < 0.05$ ), and between [0–2 km] and [6–8 km] (MENT-CLD,  $P < 0.001$ ).

Condition had no effect on drink ingestion.

## Weight loss (► **Table 1**)

As expected, participants were lighter at the end of the run ( $P < 0.001$ ), but conditions had no effect on weight loss or on weight loss percentage.

## Discussion

Our main finding points out that the use of menthol on skin in men enhances aerobic performance in a hot and humid climate, and that there is no difference in performance gain between menthol and traditional percooling strategies. Core temperature was not raised by a faster pace, and menthol brought a lower thermal sensation. However, combining both menthol and percooling did not bring any further improvements.

First, overall performance was positively impacted by cold and menthol conditions, whereas Barwood reported no difference between menthol and water sprays (at ambient temperature) in run speed during a 5-km run held in a hot and humid environment [16]. In a laboratory-controlled study, only an acute application of an 8% menthol gel on the face increased the time limit in a high-intensity exercise by more than 20% in time-to-exhaustion tests on an ergometer [13], demonstrating a positive effect of menthol on performance.

Second, run speed decreased continuously from the start in the CTL condition (► **Fig. 1a and b**), which did not happen in the other conditions. In the study by Barwood et al. [16], there was no difference in performance between conditions, and athletes' speed remained steady during the 5-km run, whereas we found a negative effect of time on performance during our longer run. Moreover, HR (► **Fig. 1c and d**) and core body temperature (► **Fig. 2c and d**) remained stable between conditions (► **Fig. 1c and d**), even if athletes were performing better, which was also reported in other works [14, 16, 26, 29, 30]. As for the heat-related perception measures,

participants felt the same lower TS at the end of the run as reported in other works [13, 16, 26, 27, 31], whereas TC was also decreased [32] or maintained/augmented [13, 16, 26, 27] (► **Table 1**). It is confirmed here by a better acceptability of the heat both in cold and menthol conditions. RPE was not modified by menthol, meaning that subjects performed all the tests with the same maximal perceived intensity, although run duration was better for some conditions. This therefore suggests an existing performance-enhancing effect in these conditions [31]. The underlying physiological mechanisms are still to be fully understood, but the type of activity (running, cycling) or the environmental conditions (dry or humid heat) might play a role.

In our study, cold water cooling and 4% menthol produced a similar effect on performance: the first strategy has long been known to buffer and help dissipate excess metabolic heat [6], whereas the second modifies perception of environmental heat and humidity [33]. Our study design also allowed us to study the effects of separate (CLD, Ment-Amb) and combined (Ment-CLD) conditions, but we found no further improvement of the latter compared to CLD or Ment-Amb taken separately, because an additive effect could have been expected. This underlines a limit in the combination of multiple strategies to maintain thermal homeostasis when exercising in a tropical climate. If the effect of external cooling is well understood, mainly by augmenting the heat capacity, the mechanisms elicited by the internal or external use of menthol remain to be fully understood.

The absence of difference in  $T_{co}$  between conditions (► **Fig. 2c**) implies that, at a faster pace, the thermoregulatory system would have to dissipate a greater amount of metabolic heat. If we assume that the low-temperature solutions ( $\sim 6^\circ\text{C}$ ), i. e., soaking the T-shirt in CLD and MENT-CLD conditions, were able to absorb a portion of the excess heat, this mechanism was not possible in the MENT-Amb condition because the solution temperature remained above  $28^\circ\text{C}$ . Regarding the non-modified  $T_{co}$  in all conditions, this raises questions about the function of thermoregulatory mechanisms already impaired by the humid atmosphere and by the limited cooling by air convection during running [2]. Moreover, this excess of heat could not be evacuated through the process of water ingested / loss of water (sweat), because on the one hand there was no difference between the environment and beverage temperatures (heat capacity), and on the other hand, between weight loss in the different conditions (physically evacuated heat through sweat). If aerobic performance strongly depends on core temperature and by its capacity to dissipate excess heat, an alternative mechanism may play a key role in thermoregulation under these harsh conditions, here potentially triggered by menthol applied to the skin. This mechanism could be related to the effect of menthol on peripheral blood circulation. An initial study reported skin vasoconstriction when a 3.5% menthol gel was applied to the skin [34], which was noted only during the first 5 min after application. In more recent studies, a vasodilatation phenomenon was observed from 5 min after application and was maximal between 15 and 45 min. A dose-dependent activation was also recorded, optimal from a 4–7% concentration [35], and was associated with a proportionally related cold sensation [36]. These menthol-related mechanisms in peripheral vasodilatation would activate, in addition to TRPM8 thermoreceptors [11], multiple vasodilator pathways, such as nitric

■ Proof copy for correction only. All forms of publication, duplication or distribution prohibited under copyright law. ■

oxide (NO) and endothelium-derived hyperpolarizing factor (EDHF), hence increasing blood flow. Although these studies were performed at rest in thermoneutral conditions, these recent advances could lead to further studies in order to assess these effects during exercise in a hot/humid environment. Such studies could also shed a novel light on the central governor model [37], regulating afferent and efferent mechanisms to prevent the occurrence of bodily harm, for example by adjusting race pace under a high thermal stress. Whereas the cold sensation elicited by the use of menthol would act on the central governor through thermosensitive afferences and therefore “trick” it into adopting a higher exercise intensity [38], locally NO-mediated and EDHF actions would promote a cutaneous vasodilatation to counteract a greater metabolic heat production. This would lead to a higher core temperature, and therefore maintain system homeostasis as long as possible, or as long as the central command is under the influence of TRPM8 afferences. Regarding core temperature, it is noticeable that numerous previous studies using menthol reported no increase in  $T_{co}$  [14, 16, 26, 29, 30]. This could be interpreted as a defense mechanism, peripherally triggered, and would compensate a dysfunction from the central command potentially leading to bodily harm. Moreover, although experimental conditions were designed to preserve a single-blind protocol, it was possible that the aroma from menthol impregnated in the T-shirt fabric would reach the respiratory system through the nasal passages. As inhaled menthol provokes a large increase of ventilation at rest and exercise [39], an additional heat loss could be made through the augmented respiratory process [40].

Finally, because menthol promotes a higher exercise intensity through sensory nerve-dependent mechanisms [38], our TS data confirm a significant cooling sensation felt by the athlete after the run, as observed in other works [13, 16, 26, 27, 33, 34], potentially acting on central command. This is also confirmed in our study by a higher TA in the MENT-Amb condition (vs. CTL,  $P < 0.05$ , ► **Table 1**). However, we did not observe changes in TC (► **Table 1**), unlike other works [16, 26, 27, 34]. To summarize, when using menthol in tropical climate, athletes ran faster and felt “comfortably fresher” than in the CTL condition.

In conclusion, our work shows improved 10-km performance in ecological conditions when using a 4% menthol skin application. This enhancement was the same as using cold water or cold menthol. The use of a higher menthol concentration (4%) than those used in most studies underlines a dose effect of menthol, whereas physiological parameters, such body core temperature and heart rate, did not show further impaired thermal stress compared to control condition. The underlying mechanisms are not yet fully understood, but recent findings on menthol-related cutaneous vasodilatation open new perspectives of research and shed a novel light on the central governor theory. Regarding the potential physiological mechanisms induced by the application of menthol on skin, future studies may focus on the effects of different concentrations of menthol on aerobic performance, especially in long-duration running.

## Funding

This work was supported by the « Programme Opérationnel - Fonds Européen de Développement Régional » (PO-FEDER) under grant 2015-FED-213, GP0008037.

## Conflict of Interest

None

## References

- [1] Hue O. The challenge of performing aerobic exercise in tropical environments: Applied knowledge and perspectives. *Int J Sports Physiol Perform* 2011; 6: 443–454
- [2] Kenefick RW, Cheuvront SN, Sawka MN. Thermoregulatory function during the marathon. *Sports Med* 2007; 37: 312–315
- [3] Brotherhood JR. Heat stress and strain in exercise and sport. *J Sci Med Sport* 2008; 11: 6–19
- [4] Ely MR, Cheuvront SN, Roberts WO et al. Impact of weather on marathon-running performance. *Med Sci Sports Exerc* 2007; 39: 487–493
- [5] Tyler CJ. Heat-stress exercise and cooling. In: Périard JD, Racinais S Eds. *Heat Stress in Sport and Exercise: Thermophysiology of Health and Performance*. Cham: Springer International Publishing; 2019: 139–157. doi:10.1007/978-3-319-93515-7\_7
- [6] Bongers CC, Thijssen DH, Veltmeijer MT et al. Precooling and percooling (cooling during exercise) both improve performance in the heat: A meta-analytical review. *Br J Sports Med* 2015; 49: 377–384
- [7] Arngrímsson SÁ, Petitt DS, Stueck MG et al. Cooling vest worn during active warm-up improves 5-km run performance in the heat. *J Appl Physiol* 2004; 96: 1867–1874
- [8] Luomala MJ, Oksa J, Salmi JA et al. Adding a cooling vest during cycling improves performance in warm and humid conditions. *J Therm Biol* 2012; 37: 47–55
- [9] Hasegawa H, Takatori T, Komura T et al. Combined effects of pre-cooling and water ingestion on thermoregulation and physical capacity during exercise in a hot environment. *J Sports Sci* 2006; 24: 3–9
- [10] Eccles R. Menthol and related cooling compounds. *J Pharm Pharmacol* 1994; 46: 618–630
- [11] Patel T, Ishiui Y, Yosipovitch G. Menthol: A refreshing look at this ancient compound. *J Am Acad Dermatol* 2007; 57: 873–878
- [12] Stevens CJ, Taylor L, Dascombe BJ. Cooling during exercise: An overlooked strategy for enhancing endurance performance in the heat. *Sports Med* 2017; 47: 829–841
- [13] Schlader ZJ, Simmons SE, Stannard SR et al. The independent roles of temperature and thermal perception in the control of human thermoregulatory behavior. *Physiol Behav* 2011; 103: 217–224
- [14] Stevens CJ, Thoseby B, Sculley DV et al. Running performance and thermal sensation in the heat are improved with menthol mouth rinse but not ice slurry ingestion. *Scand J Med Sci Sports* 2016; 26: 1209–1216
- [15] Bright FM, Chaseling GK, Jay O et al. Self-paced exercise performance in the heat with neck cooling, menthol application, and abdominal cooling. *J Sci Med Sport* 2019; 22: 371–377
- [16] Barwood MJ, Corbett J, White DK. Spraying with 0.20% L-menthol does not enhance 5 km running performance in the heat in untrained runners. *J Sports Med Phys Fitness* 2014; 54: 595–604

- [17] De Pauw K, Roelands B, Cheung SS et al. Guidelines to classify subject groups in sport-science research. *Int J Sports Physiol Perform* 2013; 8: 111–122
- [18] Léger L, Boucher R. An indirect continuous running multistage field test: The Université de Montréal track test. *Can J Appl Sport Sci* 1980; 5: 77–84
- [19] Harriss D, Macsween A, Atkinson G. Standards for ethics in sport and exercise science research: 2018 update. *Int J Sports Med* 2017; 38: 1126–1131
- [20] Hermand E, Cassirame J, Ennequin G et al. Validation of a photoplethysmographic heart rate monitor: Polar OH1. *Int J Sports Med* 2019; 40: 462–467
- [21] Plews DJ, Scott B, Altini M et al. Comparison of heart-rate variability recording with smartphone photoplethysmography, Polar H7 chest strap, and electrocardiography. *Int J Sports Physiol Perform* 2017; 12: 1324–1328
- [22] Hardy CJ, Rejeski WJ. Not what, but how one feels: The measurement of affect during exercise. *J Sport Exerc Psych* 1989; 11: 304–317
- [23] Fanger PO. Assessment of man's thermal comfort in practice. *Br J Ind Med* 1973; 30: 313–324
- [24] Goto T, Toftum J, de Dear R et al. Thermal sensation and thermophysiological responses to metabolic step-changes. *Int J Biometeorol* 2006; 50: 323–332
- [25] Zhang Y, Zhao R. Overall thermal sensation, acceptability and comfort. *Build Environ* 2008; 43: 44–50
- [26] Barwood MJ, Corbett J, Thomas K et al. Relieving thermal discomfort: Effects of sprayed L-menthol on perception, performance, and time trial cycling in the heat. *Scand J Med Sci Sports* 2015; 25: 211–218
- [27] Barwood MJ, Corbett J, White D et al. Early change in thermal perception is not a driver of anticipatory exercise pacing in the heat. *Br J Sports Med* 2012; 46: 936–942
- [28] Xiong J, Lian Z, Zhou X et al. Effects of temperature steps on human health and thermal comfort. *Build Environ* 2015; 94: 144–154
- [29] Mündel T, Jones DA. The effects of swilling an L(-)-menthol solution during exercise in the heat. *Eur J Appl Physiol* 2010; 109: 59–65
- [30] Gillis DJ, House JR, Tipton MJ. The influence of menthol on thermoregulation and perception during exercise in warm, humid conditions. *Eur J Appl Physiol* 2010; 110: 609–618
- [31] Gillis DJ, Barwood MJ, Newton PS et al. The influence of a menthol and ethanol soaked garment on human temperature regulation and perception during exercise and rest in warm, humid conditions. *J Therm Biol* 2016; 58: 99–105
- [32] Flood TR, Waldron M, Jeffries O. Oral l-menthol reduces thermal sensation, increases work-rate and extends time to exhaustion, in the heat at a fixed rating of perceived exertion. *Eur J Appl Physiol* 2017; 117: 1501–1512
- [33] Stevens CJ, Best R. Menthol: A fresh ergogenic aid for athletic performance. *Sports Med* 2017; 47: 1035–1042
- [34] Topp R, Winchester L, Mink AM et al. Comparison of the effects of ice and 3.5 % menthol gel on blood flow and muscle strength of the lower arm. *J Sport Rehabil* 2011; 20: 355–366
- [35] Craighead DH, McCartney NB, Tumlinson JH et al. Mechanisms and time course of menthol-induced cutaneous vasodilation. *Microvasc Res* 2017; 110: 43–47
- [36] Green BG. The sensory effects of l-menthol on human skin. *Somatosens Mot Res* 1992; 9: 235–244
- [37] Noakes TD, Peltonen JE, Rusko HK. Evidence that a central governor regulates exercise performance during acute hypoxia and hyperoxia. *J Exp Biol* 2001; 204: 3225–3234
- [38] Flood TR. Menthol use for performance in hot environments. *Curr Sports Med Rep* 2018; 17: 135–139
- [39] Meamarbashi A, Rajabi A. The effects of peppermint on exercise performance. *J Int Soc Sports Nutr* 2013; 10: 15
- [40] Cain JB, Livingstone SD, Nolan RW et al. Respiratory heat loss during work at various ambient temperatures. *Respir Physiol* 1990; 79: 145–150